



## FGD3 rabbit pAb

Catalog No	BYab-08634
Isotype	IgG
Reactivity	Human; Mouse
Applications	WB
Gene Name	FGD3 ZFYVE5
Protein Name	FGD3
Immunogen	Synthesized peptide derived from human FGD3 AA range: 69-119
Specificity	This antibody detects endogenous levels of FGD3 at Human/Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1: 500-2000
Concentration	1 mg/ml
Purity	≥90%
Purity Storage Stability	≥90% -20°C/1 year
-	
Storage Stability	
Storage Stability Synonyms	
Storage Stability Synonyms Observed Band	-20°C/1 year
Storage Stability Synonyms Observed Band Cell Pathway	-20°C/1 year
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	-20°C/1 year  Cytoplasm . Cytoplasm, cytoskeleton .  function:Promotes the formation of filopodia. May activate CDC42, a member of the Ras-like family of Rho- and Rac proteins, by exchanging bound GDP for free GTP. Plays a role in regulating the actin cytoskeleton and cell shape.,similarity:Contains 1 DH (DBL-homology) domain.,similarity:Contains 1 FYVE-type zinc finger.,similarity:Contains 1 PH domain.,similarity:Contains 2 PH
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	-20°C/1 year  Cytoplasm . Cytoplasm, cytoskeleton .  function:Promotes the formation of filopodia. May activate CDC42, a member of the Ras-like family of Rho- and Rac proteins, by exchanging bound GDP for free GTP. Plays a role in regulating the actin cytoskeleton and cell shape.,similarity:Contains 1 DH (DBL-homology) domain.,similarity:Contains 1 FYVE-type zinc finger.,similarity:Contains 1 PH domain.,similarity:Contains 2 PH

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658






**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.





Western blot analysis of lysates from A431 cells, primary antibody was diluted at 1:1000, 4° over night

Nanjing BYabscience technology Co.,Ltd