



FLNC rabbit pAb

Catalog No	BYab-08676
Isotype	IgG
Reactivity	Human; Mouse
Applications	IHC;IF
Gene Name	FLNC ABPL FLN2
Protein Name	FLNC
Immunogen	Synthesized peptide derived from human FLNC AA range: 335-385
Specificity	This antibody detects endogenous levels of FLNC at Human/Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	IHC-p 1: 50-200. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	
Cell Pathway	Cytoplasm . Membrane ; Peripheral membrane protein . Cytoplasm, cytoskeleton . Cytoplasm, myofibril, sarcomere, Z line . A small amount localizes at membranes. In striated muscle cells, it predominantly localizes in myofibrillar Z lines, while a minor fraction localizes with subsarcolemme. Targeting to developing and mature Z lines is mediated by the intradomain insert.
Tissue Specificity	Highly expressed in striated muscles. Weakly expressed in thyroid, fetal brain, fetal lung, retina, spinal cord and bone marrow. Not expressed in testis, pancreas, adrenal gland, placenta, liver and kidney.
Function	developmental stage:Expressed in both differentiating and adult muscles.,disease:Defects in FLNC are the cause of autosomal dominant filaminopathy [MIM:609524, 601419]. Myofibrillar myopathy (MFM) is a neuromuscular disorder, usually with an adult onset, characterized by focal myofibrillar destruction and pathological cytoplasmic protein aggregations. Autosomal dominant filaminopathy is a form of MFM characterized by morphological features of MFM and clinical features of a limb-girdle myopathy. A

Nanjing BYabscience technology Co.,Ltd



heterozygous nonsense mutation which segregates with the disease, has been identified in the FLNC gene.,domain:Comprised of a NH2-terminal actin-binding domain, 24 internally homologous repeats and two hinge regions. Repeat 24 and the second hinge domain are important for dimer formation.,domain:The filamin 20 repeat mediates interaction with XIRP1.,domain:The intradomain insert is specific to F

Background

This gene encodes one of three related filamin genes, specifically gamma filamin. These filamin proteins crosslink actin filaments into orthogonal networks in cortical cytoplasm and participate in the anchoring of membrane proteins for the actin cytoskeleton. Three functional domains exist in filamin: an N-terminal filamentous actin-binding domain, a C-terminal self-association domain, and a membrane glycoprotein-binding domain. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Tris-EDTA,pH9.0 was used for antigen retrieval. 2 Antibody was diluted at 1:200(4° overnight).3,Secondary antibody was diluted at 1:200(room temperature, 45min).