



DGAT1 rabbit pAb

Catalog No	BYab-08717
Isotype	IgG
Reactivity	Human; Mouse;Rat
Applications	WB
Gene Name	DGAT1 AGRP1 DGAT
Protein Name	DGAT1
Immunogen	Synthesized peptide derived from human DGAT1 AA range: 405-455
Specificity	This antibody detects endogenous levels of DGAT1 at Human/Mouse/Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1: 500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	
Cell Pathway	Endoplasmic reticulum membrane ; Multi-pass membrane protein .
Tissue Specificity	
Function	catalytic activity:Acyl-CoA + 1,2-diacylglycerol = CoA + triacylglycerol.,function:Catalyzes the terminal and only committed step in triacylglycerol synthesis by using diacylglycerol and fatty acyl CoA as substrates. In contrast to DGAT2 it is not essential for survival. May be involved in VLDL (very low density lipoprotein) assembly.,pathway:Lipid metabolism; glycerolipid metabolism.,similarity:Belongs to the membrane-bound acyltransferase family. Sterol o-acyltransferase subfamily.,
Background	This gene encodes an multipass transmembrane protein that functions as a key metabolic enzyme. The encoded protein catalyzes the conversion of diacylglycerol and fatty acyl CoA to triacylglycerol. This enzyme can also transfer acyl CoA to retinol. Activity of this protein may be associated with obesity and

Nanjing BYabscience technology Co.,Ltd



other metabolic diseases. [provided by RefSeq, Jul 2013],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from HEK293 cells, primary antibody was diluted at 1:1000, 4° over night