




SSRB rabbit pAb

Catalog No BYab-08771 Isotype IgG Reactivity Human; Mouse Applications WB Gene Name SSR2 TRAPB HSD25 Protein Name SSRB Immunogen Synthesized peptide derived from human SSRB AA range: 107-157 Specificity This antibody detects endogenous levels of SSRB at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Endoplasmic reticulum membrane; Single-pass type I membrane protein. Tissue Specificity Function function:TRAP proteins are part of a complex whose function is to bind calcium to the ER membrane and thereby regulate the retention of ER resident proteins, similarity Belongs to the TRAP-beta family, subunit-Heterotetramer of TRAP-patha, TRAP-delta and TRAP-delta and TRAP-gamma. Interacts with TMEM173/STING. Background		
Reactivity	Catalog No	BYab-08771
Applications Gene Name SSR2 TRAPB HSD25 Protein Name SSRB Immunogen Synthesized peptide derived from human SSRB AA range: 107-157 Specificity This antibody detects endogenous levels of SSRB at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit.lgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity 290% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Endoplasmic reticulum membrane; Single-pass type I membrane protein. Tissue Specificity Function function:TRAP proteins are part of a complex whose function is to bind calcium to the ER membrane and thereby regulate the retention of ER resident proteins, similarity-Belongs to the TRAP-beta family, subunit Heterotetramer of TRAP-alpha, TRAP-beta, TRAP-deita and TRAP-gamma. Interacts with TMEM173/STING., Background The signal sequence receptor (SSR) is a glycosylated endoplasmic reticulum (ER) membrane receptor associated with protein franslocation across the ER membrane receptor associated with protein in ranslocation across the ER membrane. The SSR consists of 2 subunits, a 34-4D glycoprotein (alpha-SSR or SSR2). The human beta-signal sequence receptor (els SSR2) maps to chromosome bands 1/q21-q23.	Isotype	IgG
Gene Name SSR2 TRAPB HSD25 Protein Name SSRB Immunogen Synthesized peptide derived from human SSRB AA range: 107-157 Specificity This antibody detects endogenous levels of SSRB at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Endoplasmic reticulum membrane; Single-pass type I membrane protein. Function function:TRAP proteins are part of a complex whose function is to bind calcium to the ER membrane and thereby regulate the retention of ER resident proteins, similarity.Belongs to the TRAP-beta family, subunit: Heterotetramer of TRAP-alpha, TRAP-alpha, TRAP-delta and TRAP-gamma. Interacts with TMEM173/STING. Background The signal sequence receptor (SSR) is a glycosylated endoplasmic reticulum (ER) membrane. The SSR consists of 2 subunits, a 34-kD glycoprotein (alpha-SSR or SSR2). The human beta-signal sequence receptor gene (SSR2) maps to chromosome bands 1q21-q23.	Reactivity	Human; Mouse
Protein Name SSRB Immunogen Synthesized peptide derived from human SSRB AA range: 107-157 Specificity This antibody detects endogenous levels of SSRB at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Cell Pathway Cell Pathway Endoplasmic reticulum membrane; Single-pass type I membrane protein. Tissue Specificity function:TRAP proteins are part of a complex whose function is to bind calcium to the ER membrane and thereby regulate the retention of ER resident proteins, similarity:Belongs to the TRAP-beta family, subunit:Heterotetramer of TRAP-alpha, TRAP-beta, TRAP-delta and TRAP-gamma. Interacts with TMEMT/3/STING. Background The signal sequence receptor (SSR) is a glycosylated endoplasmic reticulum (ER) membrane. The SSR consists of 2 subunits, a 34-kt Dylcoprotein (alpha-SSR or SSR1) and a 22-kD glycoprotein (beta-SSR or SSR2). The human beta-signal sequence receptor gene (SSR2) maps to chromosome bands 1q21-q23.	Applications	WB
Immunogen Synthesized peptide derived from human SSRB AA range: 107-157 Specificity This antibody detects endogenous levels of SSRB at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Endoplasmic reticulum membrane; Single-pass type I membrane protein. Tissue Specificity function:TRAP proteins are part of a complex whose function is to bind calcium to the ER membrane and thereby regulate the retention of ER resident proteins. similarity.Belongs to the TRAP-beta family, subunit:Heterotetramer of TRAP-alpha, TRAP-beta, TRAP-delta and TRAP-gamma. Interacts with TMEMT/3/STING. Background The signal sequence receptor (SSR) is a glycosylated endoplasmic reticulum (ER) membrane. The SSR consists of 2 subunits, a 34-kB glycoprotein (alpha-SSR or SSR) and a 22-kD glycoprotein (beta-SSR or SSR2). The human beta-signal sequence receptor gene (SSR2) maps to chromosome bands 1q21-q23.	Gene Name	SSR2 TRAPB HSD25
Specificity This antibody detects endogenous levels of SSRB at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Endoplasmic reticulum membrane; Single-pass type I membrane protein. Tissue Specificity function:TRAP proteins are part of a complex whose function is to bind calcium to the ER membrane and thereby regulate the retention of ER resident proteins, similarity:Belongs to the TRAP-beta family, subunit:Heterotetramer of TRAP-alpha, TRAP-beta, TRAP-delta and TRAP-gamma. Interacts with TMEM173/STING. Background The signal sequence receptor (SSR) is a glycosylated endoplasmic reticulum (ER) membrane. The SSR consists of 2 subunits, a 34-kD glycoprotein (alpha-SSR or SSR2). The human beta-signal sequence receptor gene (SSR2) maps to chromosome bands 1q21-q23.	Protein Name	SSRB
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Endoplasmic reticulum membrane; Single-pass type I membrane protein. Tissue Specificity Function function:TRAP proteins are part of a complex whose function is to bind calcium to the ER membrane and thereby regulate the retention of ER resident proteins, similarity:Belongs to the TRAP-beta family, subunit:Heterotetramer of TRAP-alpha, TRAP-alpha, TRAP-delta and TRAP-gamma. Interacts with TMEM173/STING., Background The signal sequence receptor (SSR) is a glycosylated endoplasmic reticulum (ER) membrane. The SSR consists of 2 subunits, a 34-kD glycoprotein (alpha-SSR or SSR1) and a 22-kD glycoprotein (beta-SSR or SSR2). The human beta-signal sequence receptor qeen (SSR2) maps to chromosome bands 1q21-q23.	Immunogen	Synthesized peptide derived from human SSRB AA range: 107-157
Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Cell Pathway Endoplasmic reticulum membrane; Single-pass type I membrane protein. Tissue Specificity Function function:TRAP proteins are part of a complex whose function is to bind calcium to the ER membrane and thereby regulate the retention of ER resident proteins, similarity:Belongs to the TRAP-beta family, subunit:Heterotetramer of TRAP-alpha, TRAP-beta, TRAP-delta and TRAP-gamma. Interacts with TMEM173/STING., Background The signal sequence receptor (SSR) is a glycosylated endoplasmic reticulum (ER) membrane receptor associated with protein translocation across the ER membrane. The SSR consists of 2 subunits, a 34-kD glycoprotein (alpha-SSR or SSR1) and a 22-kD glycoprotein (beta-SSR or SSR2). The human beta-signal sequence receptor gene (SSR2) maps to chromosome bands 1q21-q23.	Specificity	This antibody detects endogenous levels of SSRB at Human/Mouse
Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Endoplasmic reticulum membrane; Single-pass type I membrane protein. Tissue Specificity Function function:TRAP proteins are part of a complex whose function is to bind calcium to the ER membrane and thereby regulate the retention of ER resident proteins. similarity:Belongs to the TRAP-beta family.,subunit:Heterotetramer of TRAP-alpha, TRAP-beta, TRAP-delta and TRAP-gamma. Interacts with TMEM173/STING., Background The signal sequence receptor (SSR) is a glycosylated endoplasmic reticulum (ER) membrane. The SSR consists of 2 subunits, a 34-kD glycoprotein (alpha-SSR or SSR1) and a 22-kD glycoprotein (beta-SSR or SSR2). The human beta-signal sequence receptor gene (SSR2) maps to chromosome bands 1q21-q23.	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Endoplasmic reticulum membrane; Single-pass type I membrane protein. Tissue Specificity Function function:TRAP proteins are part of a complex whose function is to bind calcium to the ER membrane and thereby regulate the retention of ER resident proteins, similarity:Belongs to the TRAP-beta family, subunit:Heterotetramer of TRAP-alpha, TRAP-beta, TRAP-delta and TRAP-gamma. Interacts with TMEM173/STING., Background The signal sequence receptor (SSR) is a glycosylated endoplasmic reticulum (ER) membrane. The SSR consists of 2 subunits, a 34+kD glycoprotein (alpha-SSR or SSR1) and a 12-SkD glycoprotein (beta-SSR) or SSR2). The human beta-signal sequence receptor gene (SSR2) maps to chromosome bands 1q21-q23.	Source	Polyclonal, Rabbit,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms -20°C/1 year Observed Band Endoplasmic reticulum membrane; Single-pass type I membrane protein. Tissue Specificity Function function:TRAP proteins are part of a complex whose function is to bind calcium to the ER membrane and thereby regulate the retention of ER resident proteins., similarity:Belongs to the TRAP-beta family., subunit:Heterotetramer of TRAP-alpha, TRAP-beta, TRAP-delta and TRAP-gamma. Interacts with TMEM173/STING. Background The signal sequence receptor (SSR) is a glycosylated endoplasmic reticulum (ER) membrane receptor associated with protein translocation across the ER membrane. The SSR consists of 2 subunits, a 34-kD glycoprotein (alpha-SSR or SSR1) and a 22-kD glycoprotein (beta-SSR or SSR2). The human beta-signal sequence receptor gene (SSR2) maps to chromosome bands 1q21-q23.	Purification	
Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Endoplasmic reticulum membrane; Single-pass type I membrane protein. Tissue Specificity Function function:TRAP proteins are part of a complex whose function is to bind calcium to the ER membrane and thereby regulate the retention of ER resident proteinssimilarity:Belongs to the TRAP-beta family.,subunit:Heterotetramer of TRAP-alpha, TRAP-beta, TRAP-delta and TRAP-gamma. Interacts with TMEM173/STING., Background The signal sequence receptor (SSR) is a glycosylated endoplasmic reticulum (ER) membrane. The SSR consists of 2 subunits, a 34-kD glycoprotein (alpha-SSR or SSR1) and a 22-kD glycoprotein (beta-SSR or SSR2). The human beta-signal sequence receptor gene (SSR2) maps to chromosome bands 1q21-q23.	Dilution	WB 1: 500-2000
Synonyms Observed Band Cell Pathway Endoplasmic reticulum membrane; Single-pass type I membrane protein. Tissue Specificity Function function:TRAP proteins are part of a complex whose function is to bind calcium to the ER membrane and thereby regulate the retention of ER resident proteins.,similarity:Belongs to the TRAP-beta family.,subunit:Heterotetramer of TRAP-alpha, TRAP-beta, TRAP-delta and TRAP-gamma. Interacts with TMEM173/STING., Background The signal sequence receptor (SSR) is a glycosylated endoplasmic reticulum (ER) membrane receptor associated with protein translocation across the ER membrane. The SSR consists of 2 subunits, a 34-kD glycoprotein (alpha-SSR or SSR1) and a 22-kD glycoprotein (beta-SSR or SSR2). The human beta-signal sequence receptor gene (SSR2) maps to chromosome bands 1q21-q23.	Concentration	1 mg/ml
Synonyms Observed Band Cell Pathway Endoplasmic reticulum membrane; Single-pass type I membrane protein. Tissue Specificity Function function:TRAP proteins are part of a complex whose function is to bind calcium to the ER membrane and thereby regulate the retention of ER resident proteins.,similarity:Belongs to the TRAP-beta family.,subunit:Heterotetramer of TRAP-alpha, TRAP-beta, TRAP-delta and TRAP-gamma. Interacts with TMEM173/STING., Background The signal sequence receptor (SSR) is a glycosylated endoplasmic reticulum (ER) membrane receptor associated with protein translocation across the ER membrane. The SSR consists of 2 subunits, a 34-kD glycoprotein (alpha-SSR or SSR1) and a 22-kD glycoprotein (beta-SSR or SSR2). The human beta-signal sequence receptor gene (SSR2) maps to chromosome bands 1q21-q23.	Purity	≥90%
Cell Pathway Endoplasmic reticulum membrane; Single-pass type I membrane protein. Tissue Specificity Function function:TRAP proteins are part of a complex whose function is to bind calcium to the ER membrane and thereby regulate the retention of ER resident proteins.,similarity:Belongs to the TRAP-beta family.,subunit:Heterotetramer of TRAP-alpha, TRAP-beta, TRAP-delta and TRAP-gamma. Interacts with TMEM173/STING., Background The signal sequence receptor (SSR) is a glycosylated endoplasmic reticulum (ER) membrane receptor associated with protein translocation across the ER membrane. The SSR consists of 2 subunits, a 34-kD glycoprotein (alpha-SSR or SSR1) and a 22-kD glycoprotein (beta-SSR or SSR2). The human beta-signal sequence receptor gene (SSR2) maps to chromosome bands 1q21-q23.	Storage Stability	-20°C/1 year
Cell Pathway Endoplasmic reticulum membrane; Single-pass type I membrane protein. Tissue Specificity Function function:TRAP proteins are part of a complex whose function is to bind calcium to the ER membrane and thereby regulate the retention of ER resident proteins.,similarity:Belongs to the TRAP-beta family.,subunit:Heterotetramer of TRAP-alpha, TRAP-beta, TRAP-delta and TRAP-gamma. Interacts with TMEM173/STING., Background The signal sequence receptor (SSR) is a glycosylated endoplasmic reticulum (ER) membrane receptor associated with protein translocation across the ER membrane. The SSR consists of 2 subunits, a 34-kD glycoprotein (alpha-SSR or SSR1) and a 22-kD glycoprotein (beta-SSR or SSR2). The human beta-signal sequence receptor gene (SSR2) maps to chromosome bands 1q21-q23.		
Function function:TRAP proteins are part of a complex whose function is to bind calcium to the ER membrane and thereby regulate the retention of ER resident proteins.,similarity:Belongs to the TRAP-beta family.,subunit:Heterotetramer of TRAP-alpha, TRAP-beta, TRAP-delta and TRAP-gamma. Interacts with TMEM173/STING., Background The signal sequence receptor (SSR) is a glycosylated endoplasmic reticulum (ER) membrane receptor associated with protein translocation across the ER membrane. The SSR consists of 2 subunits, a 34-kD glycoprotein (alpha-SSR or SSR1) and a 22-kD glycoprotein (beta-SSR or SSR2). The human beta-signal sequence receptor gene (SSR2) maps to chromosome bands 1q21-q23.		
function: TRAP proteins are part of a complex whose function is to bind calcium to the ER membrane and thereby regulate the retention of ER resident proteins., similarity: Belongs to the TRAP-beta family., subunit: Heterotetramer of TRAP-alpha, TRAP-beta, TRAP-delta and TRAP-gamma. Interacts with TMEM173/STING., Background The signal sequence receptor (SSR) is a glycosylated endoplasmic reticulum (ER) membrane receptor associated with protein translocation across the ER membrane. The SSR consists of 2 subunits, a 34-kD glycoprotein (alpha-SSR or SSR1) and a 22-kD glycoprotein (beta-SSR or SSR2). The human beta-signal sequence receptor gene (SSR2) maps to chromosome bands 1q21-q23.	Synonyms	
the ER membrane and thereby regulate the retention of ER resident proteins., similarity:Belongs to the TRAP-beta family., subunit:Heterotetramer of TRAP-alpha, TRAP-beta, TRAP-delta and TRAP-gamma. Interacts with TMEM173/STING., Background The signal sequence receptor (SSR) is a glycosylated endoplasmic reticulum (ER) membrane receptor associated with protein translocation across the ER membrane. The SSR consists of 2 subunits, a 34-kD glycoprotein (alpha-SSR or SSR1) and a 22-kD glycoprotein (beta-SSR or SSR2). The human beta-signal sequence receptor gene (SSR2) maps to chromosome bands 1q21-q23.	Synonyms Observed Band	Endoplasmic reticulum membrane; Single-pass type I membrane protein.
membrane receptor associated with protein translocation across the ER membrane. The SSR consists of 2 subunits, a 34-kD glycoprotein (alpha-SSR or SSR1) and a 22-kD glycoprotein (beta-SSR or SSR2). The human beta-signal sequence receptor gene (SSR2) maps to chromosome bands 1q21-q23.	Synonyms Observed Band Cell Pathway	Endoplasmic reticulum membrane; Single-pass type I membrane protein.
	Synonyms Observed Band Cell Pathway Tissue Specificity	function:TRAP proteins are part of a complex whose function is to bind calcium to the ER membrane and thereby regulate the retention of ER resident proteins.,similarity:Belongs to the TRAP-beta family.,subunit:Heterotetramer of TRAP-alpha, TRAP-beta, TRAP-delta and TRAP-gamma. Interacts with

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658







matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

