




## INCA1 rabbit pAb

Catalog No	BYab-08796
Isotype	IgG
Reactivity	Human; Mouse
Applications	WB
Gene Name	INCA1 HSD45
Protein Name	INCA1
Immunogen	Synthesized peptide derived from human INCA1 AA range: 10-60
Specificity	This antibody detects endogenous levels of INCA1 at Human/Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1: 500-2000
Concentration	1 mg/ml
	-
Purity	≥90%
Purity Storage Stability	≥90% -20°C/1 year
Storage Stability	
Storage Stability Synonyms	
Storage Stability Synonyms Observed Band	-20°C/1 year  Nucleus . Cytoplasm .
Storage Stability Synonyms Observed Band Cell Pathway	-20°C/1 year  Nucleus . Cytoplasm .  Detected in testis, and at lower levels in ovary. Detected at very low levels in testis tumors (PubMed:15159402). Down-regulated in bone marrow cells in acute myeloid and lymphoid leukemia patients as compared with normal bone marrow
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	-20°C/1 year  Nucleus . Cytoplasm .  Detected in testis, and at lower levels in ovary. Detected at very low levels in testis tumors (PubMed:15159402). Down-regulated in bone marrow cells in acute myeloid and lymphoid leukemia patients as compared with normal bone marrow cells (PubMed:21540187).  PTM:Phosphorylated when part of a complex with CCNA1 and CDK2. Strongly phosphorylated by CDK2 on its C-terminal region spanning amino acid 149-221. Less intensively phosphorylated by CDK2 on its first 75 amino acid residues.,similarity:Belongs to the INCA family.,subunit:Interacts with CCNA1. Identified in a complex with CCNA1 and CDK2.,tissue specificity:Detected in

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658








matters needing attention

Avoid repeated freezing and thawing!

**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**



Western blot analysis of lysates from HpeG2 cells, primary antibody was diluted at 1:1000, 4° over night

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658