



ROP1B rabbit pAb

Catalog No BYab-08934 Isotype IgG Reactivity Human;Rat;Mouse; Applications WB Gene Name ROPN1B Protein Name ROP1B Immunogen Synthesized peptide derived from human ROP1B AA range: 94-144 Specificity This antibody detects endogenous levels of ROP1B at Human Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium at a source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatusing specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms	
Reactivity Human;Rat;Mouse; Applications WB Gene Name ROPN1B Protein Name ROP1B Immunogen Synthesized peptide derived from human ROP1B AA range: 94-144 Specificity This antibody detects endogenous levels of ROP1B at Human Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium at Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromat using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year	
Applications WB Gene Name ROPN1B Protein Name ROP1B Immunogen Synthesized peptide derived from human ROP1B AA range: 94-144 Specificity This antibody detects endogenous levels of ROP1B at Human Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium at Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatiusing specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year	
Gene Name ROPN1B Protein Name ROP1B Immunogen Synthesized peptide derived from human ROP1B AA range: 94-144 Specificity This antibody detects endogenous levels of ROP1B at Human Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium at a source Polyclonal, Rabbit, lgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromat using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year	
Protein Name ROP1B Immunogen Synthesized peptide derived from human ROP1B AA range: 94-144 Specificity This antibody detects endogenous levels of ROP1B at Human Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium at 20 source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromated using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year	
Immunogen Synthesized peptide derived from human ROP1B AA range: 94-144 Specificity This antibody detects endogenous levels of ROP1B at Human Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium at Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromat using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year	
Specificity This antibody detects endogenous levels of ROP1B at Human Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium at Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromat using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year	
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium at Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromat using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year	4
Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatusing specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year	
Purification The antibody was affinity-purified from rabbit serum by affinity-chromate using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year	ım azide.
Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year	
Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 year	matography
Purity ≥90% Storage Stability -20°C/1 year	
Storage Stability -20°C/1 year	
Synonyms	
Observed Band	
Cell Pathway Cell projection, cilium, flagellum. In the sperm tail, found in the principal in the cytoplasmic droplet located at the distal end of the midpiece. Innof the fibrous sheath.	cipal piece and Inner surface
Tissue Specificity	
Function domain:The RIIa domain mediates interaction with AKAP3.,miscellane name 'ropporin' comes from the Japanese word 'oppo' which means 'tail'.,similarity:Belongs to the ropporin family.,similarity:Contains 1 RIIa domain.,subcellular location:In the sperm tail, found in the principal pie	ns RIIa I piece and in Iner surface of
the cytoplasmic droplet located at the distal end of the midpiece. Inner the fibrous sheath.,subunit:Homodimer. Interacts with RHPN1. May int SPA17 (By similarity). Interacts with AKAP3.,	

Nanjing BYabscience technology Co.,Ltd






matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from Hela cells, primary antibody was diluted at 1:1000, 4° over night

Nanjing BYabscience technology Co.,Ltd