



# WFKN2 rabbit pAb

<b>Catalog No</b>	BYab-09045
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human; Mouse
<b>Applications</b>	WB
<b>Gene Name</b>	WFIKKN2 GASP1 WFIKKNRP UNQ9235/PRO31996
<b>Protein Name</b>	WFKN2
<b>Immunogen</b>	Synthesized peptide derived from human WFKN2 AA range: 506-556
<b>Specificity</b>	This antibody detects endogenous levels of WFKN2 at Human/Mouse
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
<b>Dilution</b>	WB 1: 500-2000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	
<b>Observed Band</b>	
<b>Cell Pathway</b>	Secreted .
<b>Tissue Specificity</b>	Primarily expressed in ovary, testis and brain, but not in liver. In fetal tissues, it is primarily expressed in brain, skeletal muscle, thymus and kidney.
<b>Function</b>	function:Protease-inhibitor that contains multiple distinct protease inhibitor domains. Probably has serine protease- and metalloprotease-inhibitor activity. Inhibits the biological activity of mature myostatin, but not activin.,similarity:Belongs to the WFIKKN family.,similarity:Contains 1 BPTI/Kunitz inhibitor domain.,similarity:Contains 1 Ig-like C2-type (immunoglobulin-like) domain.,similarity:Contains 1 Kazal-like domain.,similarity:Contains 1 NTR domain.,similarity:Contains 1 WAP domain.,similarity:Contains 2 BPTI/Kunitz inhibitor domains.,subunit:Interacts with both mature and propeptide myostatin/MSTN.,tissue specificity:Primarily expressed in ovary, testis and brain, but not in liver. In fetal tissues, it is primarily expressed in brain, skeletal muscle, thymus and kidney.,

Nanjing BYabscience technology Co.,Ltd



**Background**

The WFIKKN1 protein contains a WAP domain, follistatin domain, immunoglobulin domain, two tandem Kunitz domains, and an NTR domain. This gene encodes a WFIKKN1-related protein which has the same domain organization as the WFIKKN1 protein. The WAP-type, follistatin type, Kunitz-type, and NTR-type protease inhibitory domains may control the action of multiple types of proteases. [provided by RefSeq, Jul 2008],


**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

**Products Images**



Western blot analysis of lysates from HeLa cells, primary antibody was diluted at 1:1000, 4° over night