



# FAF2 rabbit pAb

<b>Catalog No</b>	BYab-09055
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human; Mouse;Rat
<b>Applications</b>	WB
<b>Gene Name</b>	FAF2 ETEA KIAA0887 UBXD8 UBXN3B
<b>Protein Name</b>	FAF2
<b>Immunogen</b>	Synthesized peptide derived from human FAF2 AA range: 120-170
<b>Specificity</b>	This antibody detects endogenous levels of FAF2 at Human/Mouse/Rat
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
<b>Dilution</b>	WB 1: 500-2000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	
<b>Observed Band</b>	
<b>Cell Pathway</b>	Cytoplasm . Lipid droplet . Endoplasmic reticulum .
<b>Tissue Specificity</b>	Broadly expressed, with highest levels in brain.
<b>Function</b>	caution:It seems odd to name this protein FAS-associated factor, but not have any mention of this in the CC lines.,miscellaneous:Up-regulated in T-cells and eosinophils from patients with atopic dermatitis.,similarity:Contains 1 UBA domain.,similarity:Contains 1 UBX domain.,tissue specificity:Broadly expressed, with highest levels in brain.,
<b>Background</b>	The protein encoded by this gene is highly expressed in peripheral blood of patients with atopic dermatitis (AD), compared to normal individuals. It may play a role in regulating the resistance to apoptosis that is observed in T cells and eosinophils of AD patients. [provided by RefSeq, Jul 2008],

Nanjing BYabscience technology Co.,Ltd




**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images



Western blot analysis of lysates from HCT116 cells, primary antibody was diluted at 1:1000, 4° over night