



## ADAP2 rabbit pAb

| Catalog No  | BYab-09928   |
|---|--|
| Isotype   | IgG  |
| Reactivity  | Human;Mouse;Rat  |
| Applications  | WB   |
| Gene Name   | ADAP2 CENTA2   |
| Protein Name  | Arf-GAP with dual PH domain-containing protein 2 (Centaurin-alpha-2) (Cnt-a2)  |
| Immunogen   | Synthesized peptide derived from human ADAP2 AA range: 301-351   |
| Specificity   | This antibody detects endogenous levels of human ADAP2   |
| Formulation   | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  |
| Source  | Polyclonal, Rabbit,IgG   |
| Purification  | The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.  |
| Dilution  | WB 1:1000-2000   |
| Concentration   | 1 mg/ml  |
| Purity  | ≥90%   |
| ·   |  |
| Storage Stability   | -20°C/1 year   |
|   | -20°C/1 year   |
| Storage Stability   | -20°C/1 year   |
| Storage Stability Synonyms  | -20°C/1 year  Cytoplasm . Cell membrane . Constitutively associated with the plasma membrane. Excluded from the nucleus.   |
| Storage Stability Synonyms Observed Band  | Cytoplasm . Cell membrane . Constitutively associated with the plasma  |
| Storage Stability Synonyms Observed Band Cell Pathway                             | Cytoplasm . Cell membrane . Constitutively associated with the plasma membrane. Excluded from the nucleus.  Highly expressed in placenta, spleen, kidney, skeletal muscle and adrenal gland. Weakly expressed in thyroid, liver, heart, lung, small intestine, peripheral blood leukocytes. Not detected in spinal cord, brain, stomach, trachea, colon, lymph |
| Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity          | Cytoplasm . Cell membrane . Constitutively associated with the plasma membrane. Excluded from the nucleus.  Highly expressed in placenta, spleen, kidney, skeletal muscle and adrenal gland. Weakly expressed in thyroid, liver, heart, lung, small intestine, peripheral blood leukocytes. Not detected in spinal cord, brain, stomach, trachea, colon, lymph |
| Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function | Cytoplasm . Cell membrane . Constitutively associated with the plasma membrane. Excluded from the nucleus.  Highly expressed in placenta, spleen, kidney, skeletal muscle and adrenal gland. Weakly expressed in thyroid, liver, heart, lung, small intestine, peripheral blood leukocytes. Not detected in spinal cord, brain, stomach, trachea, colon, lymph |

Nanjing BYabscience technology Co.,Ltd



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



| ı | Isaq | <b>P</b> S | ua | ae  | stic | าทร |
|---|------|------------|----|-----|------|-----|
| u | Jau  | C 3        | uч | uc. | วแง  | ノロコ |

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

| Products Images |  |  |  |  |  |  |
|-----------------|--|--|--|--|--|--|
|                 |  |  |  |  |  |  |
|                 |  |  |  |  |  |  |
|                 |  |  |  |  |  |  |
|                 |  |  |  |  |  |  |

Nanjing BYabscience technology Co.,Ltd