



SLAP2 rabbit pAb

Catalog No	BYab-10081
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	SLA2 C20orf156 SLAP2
Protein Name	Src-like-adapter 2 (Modulator of antigen receptor signaling) (MARS) (Src-like adapter protein 2) (SLAP-2)
Immunogen	Synthesized peptide derived from human SLAP2 AA range: 159-209
Specificity	This antibody detects endogenous levels of human SLAP2
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:1000-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Storage Stability Synonyms	-20°C/1 year
	-20°C/1 year
Synonyms	-20°C/1 year Cytoplasm.; [Isoform 1]: Cell membrane. Cytoplasmic vesicle. Localized to the plasma membrane and intracellular vesicles, including late endosomal vesicles.; [Isoform 2]: Cytoplasm . May be cytoplasmic and is not localized to membranes.
Synonyms Observed Band	Cytoplasm.; [Isoform 1]: Cell membrane. Cytoplasmic vesicle. Localized to the plasma membrane and intracellular vesicles, including late endosomal vesicles.;
Synonyms Observed Band Cell Pathway	Cytoplasm.; [Isoform 1]: Cell membrane. Cytoplasmic vesicle. Localized to the plasma membrane and intracellular vesicles, including late endosomal vesicles.; [Isoform 2]: Cytoplasm. May be cytoplasmic and is not localized to membranes. Predominantly expressed in immune system, with highest levels in peripheral blood leukocytes. Expressed in spleen, thymus and lymph nodes. Expressed in T-cells as well as in monocytes, and at low level in B-cells. Also detected in
Synonyms Observed Band Cell Pathway Tissue Specificity	Cytoplasm.; [Isoform 1]: Cell membrane. Cytoplasmic vesicle. Localized to the plasma membrane and intracellular vesicles, including late endosomal vesicles.; [Isoform 2]: Cytoplasm. May be cytoplasmic and is not localized to membranes. Predominantly expressed in immune system, with highest levels in peripheral blood leukocytes. Expressed in spleen, thymus and lymph nodes. Expressed in T-cells as well as in monocytes, and at low level in B-cells. Also detected in
Synonyms Observed Band Cell Pathway Tissue Specificity Function	Cytoplasm.; [Isoform 1]: Cell membrane. Cytoplasmic vesicle. Localized to the plasma membrane and intracellular vesicles, including late endosomal vesicles.; [Isoform 2]: Cytoplasm. May be cytoplasmic and is not localized to membranes. Predominantly expressed in immune system, with highest levels in peripheral blood leukocytes. Expressed in spleen, thymus and lymph nodes. Expressed in T-cells as well as in monocytes, and at low level in B-cells. Also detected in

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



ı	Isaq	P S	ua	ae	stic	าทร
u	Jau	C 3	uч	uc.	วแง	ノロコ

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images						

Nanjing BYabscience technology Co.,Ltd