



# CLCA1 rabbit pAb

<b>Catalog No</b>	BYab-10130
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Mouse
<b>Applications</b>	WB
<b>Gene Name</b>	CLCA1 CACC1
<b>Protein Name</b>	Calcium-activated chloride channel regulator 1 (Calcium-activated chloride channel family member 1) (hCLCA1) (Calcium-activated chloride channel protein 1) (CaCC-1) (hCaCC-1)
<b>Immunogen</b>	Synthesized peptide derived from human CLCA1 AA range: 192-242
<b>Specificity</b>	This antibody detects endogenous levels of human CLCA1
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
<b>Dilution</b>	WB 1:1000-2000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	
<b>Observed Band</b>	
<b>Cell Pathway</b>	Secreted, extracellular space . Cell membrane ; Peripheral membrane protein ; Extracellular side . Protein that remains attached to the plasma membrane appeared to be predominantly localized to microvilli.
<b>Tissue Specificity</b>	Highly expressed in small intestine and colon namely in intestinal basal crypt epithelia and goblet cells, and appendix. Weakly expressed in uterus, testis and kidney. Expressed in the airways epithelium of both asthmatic and healthy patients. Expressed in the bronchial epithelium, especially in mucus-producing goblet cells. Expressed in normal turbinate mucosa and nasal polyp. Expressed in.
<b>Function</b>	
<b>Background</b>	

Nanjing BYabscience technology Co.,Ltd



**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

**Products Images**