



ATG13 Rabbit Polyclonal Antibody

Catalog No	BYab-10293
Isotype	lgG
Reactivity	Human;Mouse;Rat
Applications	IHC;IF
Gene Name	ATG13 KIAA0652
Protein Name	Autophagy-related protein 13
Immunogen	Recombinant Protein of ATG13
Specificity	The antibody detects endogenous ATG13 protein
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	IHC-p 1:50-300. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	00%0/4
	-20°C/1 year
Synonyms	-20°C/1 year Autophagy-related protein 13
Synonyms Observed Band	•
• •	Autophagy-related protein 13
Observed Band	Autophagy-related protein 13 57kD Cytoplasm, cytosol . Preautophagosomal structure . Under starvation conditions, is localized to puncate structures primarily representing the isolation membrane; the isolation membrane sequesters a portion of the cytoplasm resulting in
Observed Band Cell Pathway	Autophagy-related protein 13 57kD Cytoplasm, cytosol . Preautophagosomal structure . Under starvation conditions, is localized to puncate structures primarily representing the isolation membrane; the isolation membrane sequesters a portion of the cytoplasm resulting in autophagosome formation Amygdala,Brain,
Observed Band Cell Pathway Tissue Specificity	Autophagy-related protein 13 57kD Cytoplasm, cytosol . Preautophagosomal structure . Under starvation conditions, is localized to puncate structures primarily representing the isolation membrane; the isolation membrane sequesters a portion of the cytoplasm resulting in autophagosome formation Amygdala,Brain, autophagic vacuole formation, autophagy, vacuole organization, cellular response to starvation, response to extracellular stimulus, macroautophagy, response to nutrient levels, cellular response to extracellular stimulus, cellular response to

Nanjing BYabscience technology Co.,Ltd



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询




matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Immunohistochemical analysis of paraffin-embedded human liver cancer. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).

Nanjing BYabscience technology Co.,Ltd