



## Peripherin Rabbit pAb

Catalog No	BYab-10308
Isotype	IgG
Reactivity	Human; Mouse;Rat
Applications	IHC;WB
Gene Name	PRPH NEF4 PRPH1
Protein Name	Peripherin
Immunogen	Synthesized peptide derived from human Peripherin AA range: 85-135
Specificity	This antibody detects endogenous levels of Peripherin at Human, Mouse,Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.15% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	IHC-p1:50-200 ,WB 1:1000-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	Peripherin (Neurofilament 4)
Observed Band	55kD
Cell Pathway	Cytoplasm, cytoskeleton . Cell projection, axon . Perikaryon .
Tissue Specificity	Expressed in the neurons of the outer hair cells in the organ of Corti and to a lesser extent in type I spiral ganglion cells.
Function	
Background	Peripherin(PRPH) Homo sapiens This gene encodes a cytoskeletal protein found in neurons of the peripheral nervous system. The encoded protein is a type III intermediate filament protein with homology to other cytoskeletal proteins such as desmin, and is a different protein that the peripherin found in photoreceptors. Mutations in this gene have been associated with susceptibility to amyotrophic lateral sclerosis.
matters needing attention	Avoid repeated freezing and thawing!

## Nanjing BYabscience technology Co.,Ltd

官方热线: 025-5229-8998

监督电话: 15950492658




国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询




**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**



Western blot analysis of 1)PC12 Cell, 2) HepG2 Cell, 3) Mouse Liver Tissue Lysate using Rabbit pAb diluted at 1:2,000.



Immunohistochemical analysis of paraffin-embedded human Squamous cell carcinoma of lung. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com

官方热线: 025-5229-8998