



BNP Rabbit pAb

Catalog No	BYab-10320
Isotype	IgG
Reactivity	Human
Applications	IHC
Gene Name	MLST8 GBL LST8
Protein Name	BNP
Immunogen	Synthesized peptide derived from human BNP AA range: 89-139
Specificity	This antibody detects endogenous levels of BNP at Human
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.90% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	IHC 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	Target of rapamycin complex subunit LST8 (TORC subunit LST8) (G protein beta subunit-like) (Gable) (Protein GbetaL) (Mammalian lethal with SEC13 protein 8) (mLST8)
Observed Band	37kD
Cell Pathway	Cytoplasm .
Tissue Specificity	Broadly expressed, with highest levels in skeletal muscle, heart and kidney.
Function	function:Unessential component of the TORC1 complex and essential component of the TORC2 complex. TORC1 controls cell growth in response to environmental signals, and is inactivated by nutrient limitation and environmental stress. Within TORC1, LST8 interacts directly with FRAP1 and enhances its kinase activity. In nutrient-poor conditions, stabilizes the FRAP1-RAPTOR interaction and favors RAPTOR-mediated inhibition of FRAP1 activity. TORC2 controls polarity of the actin cytoskeleton via the RAC1 pathway. TORC2 mediates phosphorylation of Akt/PKB on 'Ser-473' and phosphorylation of PKCalpha on 'Ser-657'. ,similarity:Belongs to the WD repeat LST8 family. ,similarity:Contains 7 WD repeats. ,subunit:Interacts with FRAP1, RAPTOR and RHEB. Part of the

Nanjing BYabscience technology Co.,Ltd



target of rapamycin complex 1 (TORC1) which contains LST8, FRAP1, RAPTOR and AKT1S1. TORC1 binds to and is inhibited by FKBP12-rapamycin. Part of

Background

Natriuretic peptide B(NPPB) Homo sapiens This gene is a member of the natriuretic peptide family and encodes a secreted protein which functions as a cardiac hormone. The protein undergoes two cleavage events, one within the cell and a second after secretion into the blood. The protein's biological actions include natriuresis, diuresis, vasorelaxation, inhibition of renin and aldosterone secretion, and a key role in cardiovascular homeostasis. A high concentration of this protein in the bloodstream is indicative of heart failure. The protein also acts as an antimicrobial peptide with antibacterial and antifungal activity. Mutations in this gene have been associated with postmenopausal osteoporosis.


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Immunohistochemical analysis of paraffin-embedded Human Heart Tissue using ILK1 Rabbit polyclonal antibody diluted at 1:200.