



# NDRG2 (Phospho-Thr348) rabbit pAb

<b>Catalog No</b>	BYab-10521
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human; Mouse;Rat
<b>Applications</b>	WB
<b>Gene Name</b>	NDRG2 KIAA1248 SYLD
<b>Protein Name</b>	NDRG2 (Phospho-Thr348)
<b>Immunogen</b>	Synthesized peptide derived from human NDRG2 (Phospho-Thr348)
<b>Specificity</b>	This antibody detects endogenous levels of NDRG2 (Phospho-Thr348) at Human, Mouse,Rat
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, and 0.177% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
<b>Dilution</b>	WB 1:500-2000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	Protein NDRG2 (N-myc downstream-regulated gene 2 protein) (Protein Syld709613)
<b>Observed Band</b>	40kD
<b>Cell Pathway</b>	Cytoplasm. Cytoplasm, perinuclear region. Cell projection, growth cone . In neurons, seems to concentrate at axonal growth cone. Perinuclear in neurons (By similarity) .
<b>Tissue Specificity</b>	Highly expressed in brain, heart, skeletal muscle and salivary gland, and moderately in kidney and liver. Expressed in dendritic cells, but not in other blood cells. Expression levels are low in pancreatic and liver cancer tissues; absent in meningioma. Expressed in low-grade gliomas but present at low levels in glioblastoma. Isoform 1 and isoform 2 are present in brain neurons and up-regulated in Alzheimer disease (at protein level).
<b>Function</b>	developmental stage:Specifically expressed during dendritic cell differentiation (in vitro). Expression is low in fetal brain and increases during brain postnatal development.,disease:Found in pathological brain lesions of Alzheimer disease.,disease:Not expressed or strongly down-regulated in various cancer types, such as astrocytoma, meningioma, liver cancer and pancreatic

Nanjing BYabscience technology Co.,Ltd



cancer.,function:May be involved in dendritic cell and neuron differentiation. May have anti-tumor activity.,similarity:Belongs to the NDRG family.,subcellular location:Perinuclear in neurons.,tissue specificity:Highly expressed in brain, heart, skeletal muscle and salivary gland, and moderately in kidney and liver. Expressed in dendritic cells, but not in other blood cells. Generally not expressed in tumor cell lines. Isoforms 1 and 2 are present in brain neurons and up-regulated in Alzheimer disease (at protein leve

**Background**

This gene is a member of the N-myc downregulated gene family which belongs to the alpha/beta hydrolase superfamily. The protein encoded by this gene is a cytoplasmic protein that may play a role in neurite outgrowth. This gene may be involved in glioblastoma carcinogenesis. Several alternatively spliced transcript variants of this gene have been described, but the full-length nature of some of these variants has not been determined. [provided by RefSeq, Jul 2008],

**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

**Products Images**

Western Blot analysis of various,using primary antibody at 1:1000 dilution. Secondary antibody(catalog#:RS23920) was diluted at 1:10000