



PBP Polyclonal Antibody

Catalog No	BYab-10693
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	IHC;IF;ELISA
Gene Name	PPBP CTAP3 CXCL7 SCYB7 TGB1 THBGB1
Protein Name	Platelet basic protein (PBP) (C-X-C motif chemokine 7) (Leukocyte-derived growth factor) (LDGF) (Macrophage-derived growth factor) (MDGF) (Small-inducible cytokine B7) [Cleaved into: Connective tissue
Immunogen	Synthetic peptide from human protein at AA range: 71-120
Specificity	The antibody detects endogenous PBP
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	IHC-p 1:50-200, ELISA 1:10000-20000. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	Platelet basic protein (PBP;C-X-C motif chemokine 7;Leukocyte-derived growth factor;LDGF;Macrophage-derived growth factor;MDGF;Small-inducible cytokine B7) [Cleaved into: Connective tissue-activating peptide III (CTAP-III;LA-PF4;Low-affinity platelet factor IV); TC-2; Connective tissue-activating peptide III(1-81;CTAP-III(1-81)); Beta-thromboglobulin (Beta-TG); Neutrophil-activating peptide 2(74;NAP-2(74)); Neutrophil-activating peptide 2(73;NAP-2(73)); Neutrophil-activating peptide 2 (NAP-2); TC-1; Neutrophil-activating peptide 2(1-66;NAP-2(1-66)); Neutrophil-activating peptide 2(1-63;NAP-2(1-63))]
Observed Band	
Cell Pathway	Secreted.
Tissue Specificity	Leukocyte,Peripheral blood monocyte,Platelet,

Nanjing BYabscience technology Co.,Ltd



Function

function:LA-PF4 stimulates DNA synthesis, mitosis, glycolysis, intracellular cAMP accumulation, prostaglandin E2 secretion, and synthesis of hyaluronic acid and sulfated glycosaminoglycan. It also stimulates the formation and secretion of plasminogen activator by human synovial cells. NAP-2 is a ligand for CXCR1 and CXCR2, and NAP-2, NAP-2(73), NAP-2(74), NAP-2(1-66), and most potent NAP-2(1-63) are chemoattractants and activators for neutrophils. TC-1 and TC-2 are antibacterial proteins, in vitro released from activated platelet alpha-granules. CTAP-III(1-81) is more potent than CTAP-III desensitize chemokine-induced neutrophil activation.,online information:CXCL7 entry,PTM:NAP-2(1-66) is produced by proteolytical processing, probably after secretion by leukocytes other than neutrophils.,PTM:NAP-2(73) and NAP-2(74) seem not be produced by proteolytical processing of secreted precursors

Background

The protein encoded by this gene is a platelet-derived growth factor that belongs to the CXC chemokine family. This growth factor is a potent chemoattractant and activator of neutrophils. It has been shown to stimulate various cellular processes including DNA synthesis, mitosis, glycolysis, intracellular cAMP accumulation, prostaglandin E2 secretion, and synthesis of hyaluronic acid and sulfated glycosaminoglycan. It also stimulates the formation and secretion of plasminogen activator by synovial cells. The protein also is an antimicrobial protein with bactericidal and antifungal activity. [provided by RefSeq, Nov 2014],

matters needing attention

Avoid repeated freezing and thawing!


Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.


Nanjing BYabscience technology Co.,Ltd



Products Images



Immunohistochemical analysis of paraffin-embedded human-brain, antibody was diluted at 1:200



Immunohistochemical analysis of paraffin-embedded human-brain, antibody was diluted at 1:200