



NECD rabbit pAb

Catalog No	BYab-09431
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	NDN
Protein Name	Necdin
Immunogen	Synthesized peptide derived from human NECD AA range: 101-151
Specificity	This antibody detects endogenous levels of human NECD
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:1000-2000
Concentration	1 mg/ml
Durity	≥90%
Purity	- 00 70
Storage Stability	-20°C/1 year
•	
Storage Stability	
Storage Stability Synonyms	
Storage Stability Synonyms Observed Band	-20°C/1 year Perikaryon. Nucleus. Neural perikarya, translocates to the nucleus of postmitotic
Storage Stability Synonyms Observed Band Cell Pathway	-20°C/1 year Perikaryon. Nucleus. Neural perikarya, translocates to the nucleus of postmitotic neurons and interacts with the nuclear matrix. Almost ubiquitous. Detected in fetal brain, lung, liver and kidney; in adult heart, brain, placenta, lung, liver, skeletal muscle, kidney, pancreas, spleen, thymus, prostate, testis, ovary, small intestine and colon. Not detected in peripheral blood
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	-20°C/1 year Perikaryon. Nucleus. Neural perikarya, translocates to the nucleus of postmitotic neurons and interacts with the nuclear matrix. Almost ubiquitous. Detected in fetal brain, lung, liver and kidney; in adult heart, brain, placenta, lung, liver, skeletal muscle, kidney, pancreas, spleen, thymus, prostate, testis, ovary, small intestine and colon. Not detected in peripheral blood
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	-20°C/1 year Perikaryon. Nucleus. Neural perikarya, translocates to the nucleus of postmitotic neurons and interacts with the nuclear matrix. Almost ubiquitous. Detected in fetal brain, lung, liver and kidney; in adult heart, brain, placenta, lung, liver, skeletal muscle, kidney, pancreas, spleen, thymus, prostate, testis, ovary, small intestine and colon. Not detected in peripheral blood

Nanjing BYabscience technology Co.,Ltd



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



ı	Isaq	P S	ua	ae	stic	าทร
u	Jau	C 3	uч	uc.	วแง	ノロコ

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images						

Nanjing BYabscience technology Co.,Ltd