




LAT1 rabbit pAb

Catalog No	BYab-09823
Isotype	IgG
Reactivity	Human;Mouse;Rat
Applications	WB
Gene Name	SLC7A5 CD98LC LAT1 MPE16
Protein Name	Large neutral amino acids transporter small subunit 1 (4F2 light chain) (4F2 LC) (4F2LC) (CD98 light chain) (Integral membrane protein E16) (L-type amino acid transporter 1) (hLAT1) (Solute carrier fa
Immunogen	Synthesized peptide derived from human LAT1 AA range: 145-195
Specificity	This antibody detects endogenous levels of human LAT1
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:1000-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	
Cell Pathway	Apical cell membrane; Multi-pass membrane protein. Cell membrane; Multi-pass membrane protein. Lysosome membrane; Multi-pass membrane protein. Located to the plasma membrane by SLC3A2/4F2hc (PubMed:9751058). Localized to the apical membrane of placental syncytiotrophoblastic cells (PubMed:11742812). Recruited to lysosomes by LAPTM4B (PubMed:25998567). Expressed in both luminal and abluminal membranes of brain capillary endothelial cells (By similarity).
Tissue Specificity	Detected in placenta, in the syncytiotrophoblast layer (at protein level) (PubMed:11389679). Expressed abundantly in adult lung, liver, brain, skeletal muscle, placenta, bone marrow, testis, resting lymphocytes and monocytes, and in fetal liver. Weaker expression in thymus, cornea, retina, peripheral leukocytes, spleen, kidney, colon and lymph node. During gestation, expression in the placenta was significantly stronger at full-term than at the mid-trimester stage. Also expressed in all human tumor cell lines tested and in the astrocytic process of primary astrocytic gliomas. Expressed in retinal endothelial cells and in the

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658







	intestinal epithelial cell line Caco-2.	
Function		
Background		
matters needing attention	Avoid repeated freezing and thawing!	
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.	

Products Images

Nanjing BYabscience technology Co.,Ltd