



DOCK6 rabbit pAb

Catalog No	BYab-09867
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB
Gene Name	DOCK6 KIAA1395
Protein Name	Dedicator of cytokinesis protein 6
Immunogen	Synthesized peptide derived from human DOCK6 AA range: 1460-1510
Specificity	This antibody detects endogenous levels of human DOCK6
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:1000-2000
Concentration	1 mg/ml
Concentration Purity	1 mg/ml ≥90%
	-
Purity	≥90%
Purity Storage Stability	≥90%
Purity Storage Stability Synonyms	≥90%
Purity Storage Stability Synonyms Observed Band	≥90% -20°C/1 year
Purity Storage Stability Synonyms Observed Band Cell Pathway	≥90% -20°C/1 year Cytoplasm . Cytoplasm, perinuclear region . Mainly located near the cell surface. Widely expressed. Expressed at low level in spleen, cerebellum, hippocampus
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	≥90% -20°C/1 year Cytoplasm . Cytoplasm, perinuclear region . Mainly located near the cell surface. Widely expressed. Expressed at low level in spleen, cerebellum, hippocampus
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	≥90% -20°C/1 year Cytoplasm . Cytoplasm, perinuclear region . Mainly located near the cell surface. Widely expressed. Expressed at low level in spleen, cerebellum, hippocampus
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function Background matters needing	≥90% -20°C/1 year Cytoplasm . Cytoplasm, perinuclear region . Mainly located near the cell surface. Widely expressed. Expressed at low level in spleen, cerebellum, hippocampus and in substantia nigra.

Nanjing BYabscience technology Co.,Ltd





Products Images

Nanjing BYabscience technology Co.,Ltd

网址:www.njbybio.com 官方热线:025-5229-8998 监督电话:15950492658