



# S26A5 Polyclonal Antibody

<b>Catalog No</b>	BYab-10844
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human; Mouse; Rat
<b>Applications</b>	WB;IHC;IF;ELISA
<b>Gene Name</b>	SLC26A5 PRES
<b>Protein Name</b>	S26A5
<b>Immunogen</b>	Synthesized peptide derived from human S26A5
<b>Specificity</b>	This antibody detects endogenous levels of human S26A5
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
<b>Dilution</b>	IHC-p 1:100-500. IF 1:50-200
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	Prestin (Solute carrier family 26 member 5)
<b>Observed Band</b>	81kD
<b>Cell Pathway</b>	Cell membrane ; Multi-pass membrane protein . Lateral wall of outer hair cells. .
<b>Tissue Specificity</b>	Organ of Corti,PCR rescued clones,
<b>Function</b>	disease:Defects in SLC26A5 are a cause of some forms of recessive non-syndromic deafness.,function:Motor protein that converts auditory stimuli to length changes in outer hair cells and mediates sound amplification in the mammalian hearing organ. Prestin is a bidirectional voltage-to-force converter, it can operate at microsecond rates. It uses cytoplasmic anions as extrinsic voltage sensors, probably chloride and bicarbonate. After binding to a site with millimolar affinity, these anions are translocated across the membrane in response to changes in the transmembrane voltage. They move towards the extracellular surface following hyperpolarization, and towards the cytoplasmic side in response to depolarization. As a consequence, this translocation triggers conformational changes in the protein that ultimately alter its surface area in the plane of the plasma membrane. The area decreases

Nanjing BYabscience technology Co.,Ltd



### Background

This gene encodes a member of the SLC26A/SulP transporter family. The protein functions as a molecular motor in motile outer hair cells (OHCs) of the cochlea, inducing changes in cell length that act to amplify sound levels. The transmembrane protein is an incomplete anion transporter, and does not allow anions to cross the cell membrane but instead undergoes a conformational change in response to changes in intracellular Cl<sup>-</sup> levels that results in a change in cell length. The protein functions at microsecond rates, which is several orders of magnitude faster than conventional molecular motor proteins. Mutations in this gene are potential candidates for causing neurosensory deafness. Multiple transcript variants encoding different isoforms have been found for this gene.[provided by RefSeq, Nov 2009],


### matters needing attention

Avoid repeated freezing and thawing!

### Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).