



# PMGT1 rabbit pAb

<b>Catalog No</b>	BYab-10946
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human; Mouse;Rat
<b>Applications</b>	WB;IHC
<b>Gene Name</b>	POMGNT1 MGAT1.2 UNQ746/PRO1475
<b>Protein Name</b>	PMGT1
<b>Immunogen</b>	Synthesized peptide derived from human PMGT1 AA range: 171-221
<b>Specificity</b>	This antibody detects endogenous levels of PMGT1 at Human/Mouse/Rat
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
<b>Dilution</b>	WB 1:500-2000;IHC-p 1:50-300
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	
<b>Observed Band</b>	
<b>Cell Pathway</b>	Golgi apparatus membrane ; Single-pass type II membrane protein .
<b>Tissue Specificity</b>	Constitutively expressed. An additional weaker band is also detected in spinal cord, lymph node, and trachea. Expressed especially in astrocytes. Also expressed in immature and mature neurons.
<b>Function</b>	catalytic activity:UDP-N-acetyl-D-glucosamine + Man-R = N-acetyl-D-glucosamine-beta-1,2-Man-R + UDP.,cofactor:Manganese.,disease:Defects in POMGNT1 are a cause of Walker-Warburg syndrome (WWS) [MIM:236670]; also known as hydrocephalus-agyria-retinal dysplasia or HARD syndrome. WWS is an autosomal recessive disorder characterized by cobblestone lissencephaly, hydrocephalus, agyria, retinal dysplasia, with or without encephalocele. It is often associated with congenital muscular dystrophy and usually lethal within the first few months of life.,disease:Defects in POMGNT1 are the cause of muscle-eye-brain disease (MEB) [MIM:253280]. MEB is an autosomal recessive disorder characterized by congenital muscular dystrophy, ocular abnormalities,

Nanjing BYabscience technology Co.,Ltd



cobblestone lissencephaly and cerebellar hypoplasia. MEB patients present severe congenital myopia, congenital glaucoma, pallor of the optic disks, retina

### Background

This gene encodes a type II transmembrane protein that resides in the Golgi apparatus. It participates in O-mannosyl glycosylation and is specific for alpha linked terminal mannose. Mutations in this gene may be associated with muscle-eye-brain disease and several congenital muscular dystrophies. Alternatively spliced transcript variants that encode different protein isoforms have been described. [provided by RefSeq, Feb 2014],


### matters needing attention

Avoid repeated freezing and thawing!


### Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images



Western blot analysis of lysates from 293T cells, primary antibody was diluted at 1:1000, 4° over night



Immunohistochemical analysis of paraffin-embedded human Colon cancer. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA, pH9.0 was used for antigen retrieval. 3, Secondary antibody was diluted at 1:200(room temperature, 45min).

Nanjing BYabscience technology Co.,Ltd