



# DGLB rabbit pAb

|                           |   |
|---------------------------|---|
| <b>Catalog No</b>         | BYab-10956  |
| <b>Isotype</b>            | IgG   |
| <b>Reactivity</b>         | Human; Mouse;Rat  |
| <b>Applications</b>       | WB  |
| <b>Gene Name</b>          | DAGLB   |
| <b>Protein Name</b>       | DGLB  |
| <b>Immunogen</b>          | Synthesized peptide derived from human DGLB AA range: 616-666   |
| <b>Specificity</b>        | This antibody detects endogenous levels of DGLB at Human/Mouse/Rat  |
| <b>Formulation</b>        | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.   |
| <b>Source</b>             | Polyclonal, Rabbit,IgG  |
| <b>Purification</b>       | The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.   |
| <b>Dilution</b>           | WB 1: 500-2000  |
| <b>Concentration</b>      | 1 mg/ml   |
| <b>Purity</b>             | ≥90%  |
| <b>Storage Stability</b>  | -20°C/1 year  |
| <b>Synonyms</b>           |   |
| <b>Observed Band</b>      |   |
| <b>Cell Pathway</b>       | Cell membrane ; Multi-pass membrane protein .   |
| <b>Tissue Specificity</b> |   |
| <b>Function</b>           | cofactor:Calcium.,enzyme regulation:Inhibited by p-hydroxy-mercuri-benzoate and HgCl(2), but not to PMSF. Also inhibited by RHC80267, a drug that blocks 2-AG formation.,function:Catalyzes the hydrolysis of diacylglycerol (DAG) to 2-arachidonoyl-glycerol (2-AG), the most abundant endocannabinoid in tissues. Required for axonal growth during development and for retrograde synaptic signaling at mature synapses.,similarity:Belongs to the AB hydrolase superfamily. Lipase family., |
| <b>Background</b>         |   |

Nanjing BYabscience technology Co.,Ltd



**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

**Products Images**

