



ZN449 rabbit pAb

Catalog No BYab-11091 Isotype IgG Reactivity Human;Rat;Mouse; Applications WB Gene Name ZNF449 ZSCAN19 Protein Name ZN449 Immunogen Synthesized peptide derived from human ZN449 AA range: 254-304 Specificity This antibody detects endogenous levels of ZN449 at Human Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Cell Pathway Observed Band Cell Pathway Cell Pathway Nucleus . Tissue Specificity Function Function krueppel C2H2-type zinc-finger protein family .similarity:Contains 1 SCAN box domain ,similarity:Contains 7 C2H2-type zinc fingers. Background This gene encodes a nuclear protein that likely functions as a transcription finger moltis. [provided by R		
Reactivity Human;Rat;Mouse; Applications WB Gene Name ZNF449 ZSCAN19 Protein Name ZN449 Immunogen Synthesized peptide derived from human ZN449 AA range: 254-304 Specificity This antibody detects endogenous levels of ZN449 at Human Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Nucleus . Tissue Specificity Function function:May be involved in transcriptional regulation, similarity:Belongs to the krueppel C2H2-type zinc-finger protein family, similarity:Contains 1 SCAN box domain, similarity:Contains 7 C2H2-type zinc fingers. Background This gene encodes a nuclear protein that likely functions as a transcription factor. The protein includes an N-terminal SCAN domain, and seven C2H2-type zinc finger motifs. [provided by RefSeq, May 2010], matters needing Avoid repeated freezing and thawing!	Catalog No	BYab-11091
Applications WB Gene Name ZNF449 ZSCAN19 Protein Name ZN449 Immunogen Synthesized peptide derived from human ZN449 AA range: 254-304 Specificity This antibody detects endogenous levels of ZN449 at Human Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Nucleus . Tissue Specificity Function function: May be involved in transcriptional regulation , similarity: Belongs to the krueppel C2H2-type zinc-finger protein family, similarity: Contains 1 SCAN box domain., similarity: Contains 7 C2H2-type zinc fingers., Background This gene encodes a nuclear protein that likely functions as a transcription factor. The protein includes an N-terminal SCAN domain, and seven C2H2-type zinc finger motifs. [provided by RefSeq, May 2010], Matters needing Avoid repeated freezing and thaving!	Isotype	IgG
Gene Name ZNF449 ZSCAN19 Protein Name ZN449 Immunogen Synthesized peptide derived from human ZN449 AA range: 254-304 Specificity This antibody detects endogenous levels of ZN449 at Human Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Nucleus . Tissue Specificity Function function:May be involved in transcriptional regulation, similarity:Belongs to the krueppel C2H2-type zinc-finger protein family, similarity:Contains 1 SCAN box domain, similarity:Contains 7 C2H2-type zinc fingers. Background This gene encodes a nuclear protein that likely functions as a transcription factor. The protein includes an N-terminal SCAN domain, and seven C2H2-type zinc finger motifs. [provided by RefSeq, May 2010], matters needing Avoid repeated freezing and thawing!	Reactivity	Human;Rat;Mouse;
Protein Name ZN449 Immunogen Synthesized peptide derived from human ZN449 AA range: 254-304 Specificity This antibody detects endogenous levels of ZN449 at Human Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Nucleus : Tissue Specificity Function function:May be involved in transcriptional regulation., similarity:Belongs to the krueppel C2/H2-type zinc-finger protein family, similarity:Contains 1 SCAN box domain, similarity:Contains 7 C2H2-type zinc fingers., Background This gene encodes a nuclear protein that likely functions as a transcription factor. The protein includes an N-terminal SCAN domain, and seven C2H2-type zinc finger motifs. [provided by RefSeq, May 2010], matters needing Avoid repeated freezing and thawing!	Applications	WB
Immunogen Synthesized peptide derived from human ZN449 AA range: 254-304 Specificity This antibody detects endogenous levels of ZN449 at Human Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Nucleus . Tissue Specificity Function function:May be involved in transcriptional regulation.,similarity:Belongs to the krueppel C2H2-type zinc-finger protein family, similarity:Contains 1 SCAN box domain., similarity:Contains 7 C2H2-type zinc fingers. Background This gene encodes a nuclear protein that likely functions as a transcription factor. The protein includes an N-terminal SCAN domain, and seven C2H2-type zinc finger motifs. [provided by RefSeq, May 2010], matters needing Avoid repeated freezing and thawing!	Gene Name	ZNF449 ZSCAN19
Specificity This antibody detects endogenous levels of ZN449 at Human Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Nucleus. Tissue Specificity Function function:May be involved in transcriptional regulation, similarity:Belongs to the krueppel C2H2-type zinc-finger protein family, similarity:Contains 1 SCAN box domain., similarity:Contains 7 C2H2-type zinc fingers., Background This gene encodes a nuclear protein that likely functions as a transcription factor. The protein includes an N-terminal SCAN domain, and seven C2H2-type zinc finger motifs. [provided by RefSeq, May 2010], matters needing Avoid repeated freezing and thawing!	Protein Name	ZN449
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Nucleus. Tissue Specificity Function function:May be involved in transcriptional regulation, similarity:Belongs to the krueppel C2H2-type zinc-finger protein family, similarity:Contains 1 SCAN box domain, similarity:Contains 7 C2H2-type zinc fingers., Background This gene encodes a nuclear protein that likely functions as a transcription factor. The protein includes an N-terminal SCAN domain, and seven C2H2-type zinc finger motifs. [provided by RefSeq, May 2010], matters needing Avoid repeated freezing and thawing!	Immunogen	Synthesized peptide derived from human ZN449 AA range: 254-304
Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Nucleus . Tissue Specificity Function function:May be involved in transcriptional regulation.,similarity:Belongs to the krueppel C2H2-type zinc-finger protein family ,similarity:Contains 1 SCAN box domain.,similarity:Contains 7 C2H2-type zinc fingers., Background This gene encodes a nuclear protein that likely functions as a transcription factor. The protein includes an N-terminal SCAN domain, and seven C2H2-type zinc finger motifs. [provided by RefSeq, May 2010], matters needing Avoid repeated freezing and thawing!	Specificity	This antibody detects endogenous levels of ZN449 at Human
Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Nucleus . Tissue Specificity Function function:May be involved in transcriptional regulation.,similarity:Belongs to the krueppel C2H2-type zinc-finger protein family, similarity:Contains 1 SCAN box domain.,similarity:Contains 7 C2H2-type zinc fingers., Background This gene encodes a nuclear protein that likely functions as a transcription factor. The protein includes an N-terminal SCAN domain, and seven C2H2-type zinc finger motifs. [provided by RefSeq, May 2010], matters needing Avoid repeated freezing and thawing!	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Nucleus . Tissue Specificity Function function:May be involved in transcriptional regulation.,similarity:Belongs to the krueppel C2H2-type zinc-finger protein familysimilarity:Contains 1 SCAN box domain.,similarity:Contains 7 C2H2-type zinc fingers., Background This gene encodes a nuclear protein that likely functions as a transcription factor. The protein includes an N-terminal SCAN domain, and seven C2H2-type zinc finger motifs. [provided by RefSeq, May 2010], matters needing Avoid repeated freezing and thawing!	Source	Polyclonal, Rabbit,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Nucleus . Tissue Specificity Function function:May be involved in transcriptional regulation.,similarity:Belongs to the krueppel C2H2-type zinc-finger protein family.,similarity:Contains 1 SCAN box domain.,similarity:Contains 7 C2H2-type zinc fingers., Background This gene encodes a nuclear protein that likely functions as a transcription factor. The protein includes an N-terminal SCAN domain, and seven C2H2-type zinc finger motifs. [provided by RefSeq, May 2010], matters needing Avoid repeated freezing and thawing!	Purification	
Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Nucleus . Tissue Specificity Function function:May be involved in transcriptional regulation.,similarity:Belongs to the krueppel C2H2-type zinc-finger protein family.,similarity:Contains 1 SCAN box domain.,similarity:Contains 7 C2H2-type zinc fingers., Background This gene encodes a nuclear protein that likely functions as a transcription factor. The protein includes an N-terminal SCAN domain, and seven C2H2-type zinc finger motifs. [provided by RefSeq, May 2010], matters needing Avoid repeated freezing and thawing!	Dilution	WB 1: 500-2000
Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Nucleus . Tissue Specificity Function function:May be involved in transcriptional regulation.,similarity:Belongs to the krueppel C2H2-type zinc-finger protein family.,similarity:Contains 1 SCAN box domain.,similarity:Contains 7 C2H2-type zinc fingers., Background This gene encodes a nuclear protein that likely functions as a transcription factor. The protein includes an N-terminal SCAN domain, and seven C2H2-type zinc finger motifs. [provided by RefSeq, May 2010], matters needing Avoid repeated freezing and thawing!	Concentration	1 mg/ml
Synonyms Observed Band Cell Pathway Nucleus . Tissue Specificity Function function:May be involved in transcriptional regulation.,similarity:Belongs to the krueppel C2H2-type zinc-finger protein family.,similarity:Contains 1 SCAN box domain.,similarity:Contains 7 C2H2-type zinc fingers., Background This gene encodes a nuclear protein that likely functions as a transcription factor. The protein includes an N-terminal SCAN domain, and seven C2H2-type zinc finger motifs. [provided by RefSeq, May 2010], matters needing Avoid repeated freezing and thawing!	Purity	≥90%
Observed Band Cell Pathway Nucleus . Tissue Specificity Function function:May be involved in transcriptional regulation.,similarity:Belongs to the krueppel C2H2-type zinc-finger protein family.,similarity:Contains 1 SCAN box domain.,similarity:Contains 7 C2H2-type zinc fingers., Background This gene encodes a nuclear protein that likely functions as a transcription factor. The protein includes an N-terminal SCAN domain, and seven C2H2-type zinc finger motifs. [provided by RefSeq, May 2010], matters needing Avoid repeated freezing and thawing!	Storage Stability	-20°C/1 year
Cell Pathway Nucleus . Tissue Specificity Function function:May be involved in transcriptional regulation.,similarity:Belongs to the krueppel C2H2-type zinc-finger protein family.,similarity:Contains 1 SCAN box domain.,similarity:Contains 7 C2H2-type zinc fingers., Background This gene encodes a nuclear protein that likely functions as a transcription factor. The protein includes an N-terminal SCAN domain, and seven C2H2-type zinc finger motifs. [provided by RefSeq, May 2010], matters needing Avoid repeated freezing and thawing!	Synonyms	
Tissue Specificity Function function:May be involved in transcriptional regulation.,similarity:Belongs to the krueppel C2H2-type zinc-finger protein family.,similarity:Contains 1 SCAN box domain.,similarity:Contains 7 C2H2-type zinc fingers., Background This gene encodes a nuclear protein that likely functions as a transcription factor. The protein includes an N-terminal SCAN domain, and seven C2H2-type zinc finger motifs. [provided by RefSeq, May 2010], matters needing Avoid repeated freezing and thawing!	Observed Band	
Function function:May be involved in transcriptional regulation.,similarity:Belongs to the krueppel C2H2-type zinc-finger protein family.,similarity:Contains 1 SCAN box domain.,similarity:Contains 7 C2H2-type zinc fingers., This gene encodes a nuclear protein that likely functions as a transcription factor. The protein includes an N-terminal SCAN domain, and seven C2H2-type zinc finger motifs. [provided by RefSeq, May 2010], Matters needing Avoid repeated freezing and thawing!	Cell Pathway	Nucleus .
krueppel C2H2-type zinc-finger protein family.,similarity:Contains 1 SCAN box domain.,similarity:Contains 7 C2H2-type zinc fingers., Background This gene encodes a nuclear protein that likely functions as a transcription factor. The protein includes an N-terminal SCAN domain, and seven C2H2-type zinc finger motifs. [provided by RefSeq, May 2010], matters needing Avoid repeated freezing and thawing!	Tissue Specificity	
This gene encodes a nuclear protein that likely functions as a transcription factor. The protein includes an N-terminal SCAN domain, and seven C2H2-type zinc finger motifs. [provided by RefSeq, May 2010], Matters needing Avoid repeated freezing and thawing!	Function	function:May be involved in transcriptional regulation.,similarity:Belongs to the krueppel C2H2-type zinc-finger protein family.,similarity:Contains 1 SCAN box domain.,similarity:Contains 7 C2H2-type zinc fingers.,
	Background	This gene encodes a nuclear protein that likely functions as a transcription factor. The protein includes an N-terminal SCAN domain, and seven C2H2-type zinc
	matters needing attention	Avoid repeated freezing and thawing!

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658




国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from MCF-7 cells, primary antibody was diluted at 1:1000, 4° over night

Nanjing BYabscience technology Co.,Ltd