



# URAD rabbit pAb

<b>Catalog No</b>	BYab-11172
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human; Mouse
<b>Applications</b>	WB
<b>Gene Name</b>	PRHOXNB
<b>Protein Name</b>	URAD
<b>Immunogen</b>	Synthesized peptide derived from human URAD AA range: 80-130
<b>Specificity</b>	This antibody detects endogenous levels of URAD at Human/Mouse
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
<b>Dilution</b>	WB 1: 500-2000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	
<b>Observed Band</b>	
<b>Cell Pathway</b>	Peroxisome .
<b>Tissue Specificity</b>	Apparently not expressed.
<b>Function</b>	catalytic activity:5-hydroxy-2-oxo-4-ureido-2,5-dihydro-1H-imidazole-5-carboxylate = (S)-allantoin + CO(2).,caution:Could be the product of a pseudogene. In primates the genes coding for the enzymes for the degradation of uric acid were inactivated and converted to pseudogenes.,function:Catalyzes the stereoselective decarboxylation of 2-oxo-4-hydroxy-4-carboxy-5-ureidoimidazoline (OHCU) to (S)-allantoin .,pathway:Purine metabolism; uric acid degradation; (S)-allantoin from uric acid: step 3/3.,similarity:Belongs to the OHCU decarboxylase family.,tissue specificity:Apparently not expressed.,

## Background

Nanjing BYabscience technology Co.,Ltd




**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

**Products Images**



Nanjing BYabscience technology Co.,Ltd