



# ITM2C rabbit pAb

<b>Catalog No</b>	BYab-11286
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human; Mouse;Rat
<b>Applications</b>	WB;IHC
<b>Gene Name</b>	ITM2C BRI3 hucep-14 NPD018 PSEC0047
<b>Protein Name</b>	ITM2C
<b>Immunogen</b>	Synthesized peptide derived from human ITM2C AA range: 1-51
<b>Specificity</b>	This antibody detects endogenous levels of ITM2C at Human/Mouse/Rat
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
<b>Dilution</b>	WB 1:500-2000;IHC-p 1:50-300
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	
<b>Observed Band</b>	
<b>Cell Pathway</b>	Lysosome membrane ; Single-pass type II membrane protein . Cell membrane ; Single-pass type II membrane protein .
<b>Tissue Specificity</b>	High levels in the brain, specifically in the cerebral cortex, medulla, amygdala, hippocampus, thalamus, caudate nucleus, cerebellum, olfactory lobe and spinal cord. Very low levels in other organs.
<b>Function</b>	PTM:Type I membrane-bound, as well as soluble, furin has a pre-eminent role in ITM2C proteolytic processing. PCSK7 and PCSK5 may also be involved although to a lesser extent. The soluble form of PCSK7 is incapable of processing ITM2C.,similarity:Belongs to the ITM2 family.,similarity:Contains 1 BRICHOS domain.,subunit:Interacts with BACE1.,tissue specificity:High levels in the brain, specifically in the cerebral cortex, medulla, amygdala, hippocampus, thalamus, caudate nucleus, cerebellum, olfactory lobe and spinal cord. Very low levels in other organs.,
<b>Background</b>	

Nanjing BYabscience technology Co.,Ltd




**matters needing attention**

Avoid repeated freezing and thawing!


**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images



Western blot analysis of lysates from MDA-MB cells, primary antibody was diluted at 1:1000, 4° over night



Immunohistochemical analysis of paraffin-embedded human Squamous cell carcinoma of lung. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).