



## L12R1 rabbit pAb

Catalog No	BYab-11373
Isotype	IgG
Reactivity	Human; Mouse
Applications	WB;IHC
Gene Name	LOH12CR1
Protein Name	L12R1
Immunogen	Synthesized peptide derived from human L12R1 AA range: 9-59
Specificity	This antibody detects endogenous levels of L12R1 at Human/Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification Dilution	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. WB 1:500-2000;IHC-p 1:50-300
Component and in a	1 mg/ml
Concentration	r mg/m
Concentration	≥90%
	-
Purity	≥90%
Purity Storage Stability	≥90%
Purity Storage Stability Synonyms	≥90%
Purity Storage Stability Synonyms Observed Band	≥90% -20°C/1 year
Purity Storage Stability Synonyms Observed Band Cell Pathway	≥90% -20°C/1 year Lysosome membrane ; Lipid-anchor ; Cytoplasmic side .
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	≥90% -20°C/1 year Lysosome membrane ; Lipid-anchor ; Cytoplasmic side . Ubiquitously expressed (at protein level).
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	≥90% -20°C/1 year Lysosome membrane ; Lipid-anchor ; Cytoplasmic side . Ubiquitously expressed (at protein level).
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function Background matters needing	≥90% -20°C/1 year Lysosome membrane ; Lipid-anchor ; Cytoplasmic side . Ubiquitously expressed (at protein level). similarity:Belongs to the LOH12CR1 family.,tissue specificity:Widely expressed.,


## Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658






## **Products Images**



Immunohistochemical analysis of paraffin-embedded human Squamous cell carcinoma of lung. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).

Western blot analysis of lysates from PC-12 cells,

primary antibody was diluted at 1:1000, 4°over night



Nanjing BYabscience technology Co.,Ltd