



TM9S4 rabbit pAb

Catalog No	BYab-11512
Isotype	IgG
Reactivity	Human; Mouse;Rat
Applications	WB
Gene Name	TM9SF4 KIAA0255
Protein Name	TM9S4
Immunogen	Synthesized peptide derived from human TM9S4 AA range: 315-365
Specificity	This antibody detects endogenous levels of TM9S4 at Human/Mouse/Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1: 500-2000
Concentration	1 mg/ml
Concentration Purity	1 mg/ml ≥90%
	-
Purity	≥90%
Purity Storage Stability	≥90%
Purity Storage Stability Synonyms	≥90%
Purity Storage Stability Synonyms Observed Band	≥90% -20°C/1 year
Purity Storage Stability Synonyms Observed Band Cell Pathway	≥90% -20°C/1 year Membrane; Multi-pass membrane protein. Golgi apparatus. Early endosome. Highly expressed in metastatic melanoma cells whereas it is undetectable in primary melanoma cells, healthy skin tissues and peripheral blood lymphocytes. Expressed in CD34(+) hematopoietic progenitor cells and during monocyte and granulocyte differentiation. Overexpressed in acute myeloid leukemia, in
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	≥90% -20°C/1 year Membrane ; Multi-pass membrane protein . Golgi apparatus . Early endosome . Highly expressed in metastatic melanoma cells whereas it is undetectable in primary melanoma cells, healthy skin tissues and peripheral blood lymphocytes. Expressed in CD34(+) hematopoietic progenitor cells and during monocyte and granulocyte differentiation. Overexpressed in acute myeloid leukemia, in particular in those displaying granulocytic differentiation (at protein level).
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	≥90% -20°C/1 year Membrane ; Multi-pass membrane protein . Golgi apparatus . Early endosome . Highly expressed in metastatic melanoma cells whereas it is undetectable in primary melanoma cells, healthy skin tissues and peripheral blood lymphocytes. Expressed in CD34(+) hematopoietic progenitor cells and during monocyte and granulocyte differentiation. Overexpressed in acute myeloid leukemia, in particular in those displaying granulocytic differentiation (at protein level).

Nanjing BYabscience technology Co.,Ltd




国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from HuvEc cells, primary antibody was diluted at 1:1000, 4° over night

Nanjing BYabscience technology Co.,Ltd