



OXA1L rabbit pAb

Catalog No	BYab-11597
Isotype	IgG
Reactivity	Human; Mouse
Applications	WB
Gene Name	OXA1L
Protein Name	OXA1L
Immunogen	Synthesized peptide derived from human OXA1L AA range: 53-103
Specificity	This antibody detects endogenous levels of OXA1L at Human/Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1: 500-2000
	4 manufact
Concentration	1 mg/ml
Concentration Purity	1 mg/ml ≥90%
Purity	≥90%
Purity Storage Stability	≥90%
Purity Storage Stability Synonyms	≥90%
Purity Storage Stability Synonyms Observed Band	≥90% -20°C/1 year
Purity Storage Stability Synonyms Observed Band Cell Pathway	≥90% -20°C/1 year
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	≥90% -20°C/1 year Mitochondrion inner membrane ; Multi-pass membrane protein . function:Required for the insertion of integral membrane proteins into the mitochondrial inner membrane. Essential for the activity and assembly of
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	≥90% -20°C/1 year Mitochondrion inner membrane; Multi-pass membrane protein. function:Required for the insertion of integral membrane proteins into the mitochondrial inner membrane. Essential for the activity and assembly of cytochrome oxidase.,similarity:Belongs to the OXA1/oxaA family., This gene encodes an evolutionarily conserved protein that is localized to the inner mitochondrial membrane. The encoded protein is essential for the translocation of the N-terminal tail of subunit 2 of cytochrome c oxidase, and is involved in the assembly of the cytochrome c oxidase and ATPase complexes of

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658




国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.





Western blot analysis of lysates from HpeG2 cells, primary antibody was diluted at 1:1000, 4° over night

Nanjing BYabscience technology Co.,Ltd