



# ZNT6 rabbit pAb

<b>Catalog No</b>	BYab-11651
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human; Mouse
<b>Applications</b>	WB
<b>Gene Name</b>	SLC30A6 ZNT6
<b>Protein Name</b>	ZNT6
<b>Immunogen</b>	Synthesized peptide derived from human ZNT6 AA range: 58-108
<b>Specificity</b>	This antibody detects endogenous levels of ZNT6 at Human/Mouse
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
<b>Dilution</b>	WB 1: 500-2000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	
<b>Observed Band</b>	
<b>Cell Pathway</b>	Golgi apparatus, trans-Golgi network membrane; Multi-pass membrane protein. Found in vesicles. .
<b>Tissue Specificity</b>	Expressed in brain; especially in cerebellum, hippocampus, parahippocampal gyrus, superior and middle temporal gyrus. Also expressed in B-cells, colon, eye, and lung. Lower expression was present in bone, brain, cervix, ear, heart, kidney, muscle, nerve, pancreas, prostate, skin, stomach, and testis.
<b>Function</b>	function:Zinc-efflux transporter which allocates the cytoplasmic zinc to the trans-Golgi network (TGN) as well as the vesicular compartment.,miscellaneous:Seems to have lost most of the histidine residues in the loop between the fourth and fifth transmembrane regions and appears to exert transport function by forming complexes with znt5.,similarity:Belongs to the cation diffusion facilitator (CDF) transporter (TC 2.A.4) family. SLC30A subfamily.,subcellular location:Found in vesicles.,subunit:Heterooligomer. Interacts with ZNT5.,tissue specificity:Expressed in brain; especially in cerebellum, hippocampus, parahippocampal gyrus, superior and middle temporal

Nanjing BYabscience technology Co.,Ltd



gyrus. Also expressed in B cells, colon, eye, and lung. Lower expression was present in bone, brain, cervix, ear, heart, kidney, muscle, nerve, pancreas, prostate, skin, stomach, and testis.,

**Background**

This gene encodes a member of a family of proteins that function as zinc transporters. This protein can regulate subcellular levels of zinc in the Golgi and vesicles. Expression of this gene is altered in the Alzheimer's disease brain plaques. [provided by RefSeq, Aug 2016],


**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

**Products Images**



Western blot analysis of lysates from MCF-7 cells, primary antibody was diluted at 1:1000, 4° over night