



SAC2 rabbit pAb

Catalog No	BYab-11730
Isotype	IgG
Reactivity	Human; Mouse
Applications	WB
Gene Name	INPP5F KIAA0966 SAC2 MSTP007 MSTP047
Protein Name	SAC2
Immunogen	Synthesized peptide derived from human SAC2 AA range: 1006-1056
Specificity	This antibody detects endogenous levels of SAC2 at Human/Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1: 500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	
Cell Pathway	Membrane, clathrin-coated pit . Early endosome . Recycling endosome . Also found on macropinosomes. .
Tissue Specificity	Ubiquitous (PubMed:11274189). Highly expressed in brain (PubMed:26203138).
Function	function:Inositol 5-phosphatase which acts on phosphatidylinositol 4,5-bisphosphate and phosphatidylinositol 3,4,5-triphosphate. Hydrolyzes phosphatidylinositol 4,5-bisphosphate most effectively. Modulates AKT/GSK3B pathway by decreasing AKT and GSK3B phosphorylation.,miscellaneous:Inositol polyphosphate 5-phosphatases usually have 2 conserved motifs that are essential for 5-phosphatase activity. However, INPP5F does not contain such motifs, and phosphatase activity seems to exist in the SAC domain, suggesting a new type of phosphoinositide 5-phosphatase.,similarity:Contains 1 SAC domain.,tissue specificity:Ubiquitous.,
Background	The protein encoded by this gene is an inositol 1,4,5-trisphosphate (InsP3) 5-phosphatase and contains a Sac domain. The activity of this protein is specific

Nanjing BYabscience technology Co.,Ltd



for phosphatidylinositol 4,5-bisphosphate and phosphatidylinositol 3,4,5-trisphosphate. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene. [provided by RefSeq, Aug 2011],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from DU145 cells, primary antibody was diluted at 1:1000, 4° over night