



# TDRD9 rabbit pAb

<b>Catalog No</b>	BYab-11733
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human; Mouse;Rat
<b>Applications</b>	WB
<b>Gene Name</b>	TDRD9 C14orf75
<b>Protein Name</b>	TDRD9
<b>Immunogen</b>	Synthesized peptide derived from human TDRD9 AA range: 335-385
<b>Specificity</b>	This antibody detects endogenous levels of TDRD9 at Human/Mouse/Rat
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
<b>Dilution</b>	WB 1: 500-2000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	
<b>Observed Band</b>	
<b>Cell Pathway</b>	Cytoplasm . Nucleus . Component of the nuage, also named P granule, a germ-cell-specific organelle required to repress transposon activity during meiosis. Specifically localizes to piP-bodies, a subset of the nuage which contains secondary piRNAs. PIWIL2 is required for its localization to piP-bodies. .
<b>Tissue Specificity</b>	
<b>Function</b>	similarity:Contains 1 helicase ATP-binding domain.,similarity:Contains 1 helicase C-terminal domain.,
<b>Background</b>	
<b>matters needing attention</b>	Avoid repeated freezing and thawing!


Nanjing BYabscience technology Co.,Ltd



### Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images



Western blot analysis of lysates from HpeG2 cells, primary antibody was diluted at 1:1000, 4° over night.