



RN169 rabbit pAb

IsotypeIgGReactivityHuman; MouseApplicationsWB;ELISA;IHCGene NameRNF169 KIAA1991Protein NameRN169ImmunogenSynthesized peptide derived from human RN169 AA range: 123-173SpecificityThis antibody detects endogenous levels of RN169 at Human/MouseFormulationLiquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.SourcePolyclonal, Rabbit,IgGPurificationThe antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.DilutionWB 1:500-2000;IHC-p 1:50-300; ELISA 2000-20000Concentration1 mg/mlPurify≥90%Storage Stability-20°C/1 yearSynonyms-20°C/1 yearColserved BandRNF168-dependent ubiquitin products.Tissue SpecificitySimilarity:Contains 1 RING-type zinc finger.,BackgroundAvoid repeated freezing and thawing!	Catalog No	BYab-11752
Applications WB;ELISA;IHC Gene Name RNF169 KIAA1991 Protein Name RN169 Immunogen Synthesized peptide derived from human RN169 AA range: 123-173 Specificity This antibody detects endogenous levels of RN169 at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:500-2000;IHC-p 1:50-300; ELISA 2000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Chromosome . Nucleus, nucleoplasm . Localizes to sites of double-strand breaks (DSB) following DNA damage. Recruited to DSBs via recognition of RNF168-dependent ubiquitin products Tissue Specificity similarity:Contains 1 RING-type zinc finger., Background Avoid repeated freezing and thawing!	Isotype	lgG
Gene Name RNF169 KIAA1991 Protein Name RN169 Immunogen Synthesized peptide derived from human RN169 AA range: 123-173 Specificity This antibody detects endogenous levels of RN169 at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:500-2000;IHC-p 1:50-300; ELISA 2000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms - Cell Pathway Chromosome . Nucleus, nucleoplasm . Localizes to sites of double-strand breaks (RNF168-dependent ubiquitin products Tissue Specificity similarity:Contains 1 RING-type zinc finger., Background Avoid repeated freezing and thawing!	Reactivity	Human; Mouse
Protein Name RN169 Immunogen Synthesized peptide derived from human RN169 AA range: 123-173 Specificity This antibody detects endogenous levels of RN169 at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:500-2000;IHC-p 1:50-300; ELISA 2000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms - Cell Pathway Chromosome . Nucleus, nucleoplasm . Localizes to sites of double-strand breaks (DSBs) following DNA damage. Recruited to DSBs via recognition of RNF-168-dependent ubiquitin products . Tissue Specificity - Function similarity:Contains 1 RING-type zinc finger., Background Avoid repeated freezing and thawing!	Applications	WB;ELISA;IHC
Immunogen Synthesized peptide derived from human RN169 AA range: 123-173 Specificity This antibody detects endogenous levels of RN169 at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:500-2000;IHC-p 1:50-300; ELISA 2000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Chromosome . Nucleus, nucleoplasm . Localizes to sites of double-strand breaks (DSBs) following DNA damage. Recruited to DSBs via recognition of RNF 168-dependent ubiquitin products . Tissue Specificity Function Function similarity:Contains 1 RING-type zinc finger., Background Avoid repeated freezing and thawing!	Gene Name	RNF169 KIAA1991
Specificity This antibody detects endogenous levels of RN169 at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:500-2000;IHC-p 1:50-300; ELISA 2000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Chromosome . Nucleus, nucleoplasm . Localizes to sites of double-strand breaks (DSBs) following DNA damage. Recruited to DSBs via recognition of RNF 168-dependent ubiquitin products . Tissue Specificity Function Function similarity:Contains 1 RING-type zinc finger., Background Avoid repeated freezing and thawing!	Protein Name	RN169
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:500-2000;IHC-p 1:50-300; ELISA 2000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms -20°C/1 year Observed Band Chromosome . Nucleus, nucleoplasm . Localizes to sites of double-strand breaks (DSBs) following DNA damage. Recruited to DSBs via recognition of RNF168-dependent ubiquitin products Tissue Specificity similarity:Contains 1 RING-type zinc finger., Background Avoid repeated freezing and thawing!	Immunogen	Synthesized peptide derived from human RN169 AA range: 123-173
Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:500-2000;IHC-p 1:50-300; ELISA 2000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Chromosome . Nucleus, nucleoplasm . Localizes to sites of double-strand breaks (DSBs) following DNA damage. Recruited to DSBs via recognition of RNF168-dependent ubiquitin products Tissue Specificity similarity:Contains 1 RING-type zinc finger., Background Avoid repeated freezing and thawing!	Specificity	This antibody detects endogenous levels of RN169 at Human/Mouse
Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:500-2000;IHC-p 1:50-300; ELISA 2000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms -20°C/1 year Observed Band Chromosome . Nucleus, nucleoplasm . Localizes to sites of double-strand breaks (DSBs) following DNA damage. Recruited to DSBs via recognition of RNF168-dependent ubiquitin products . Tissue Specificity similarity:Contains 1 RING-type zinc finger., Background Avoid repeated freezing and thawing!	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
using specific immunogen. Dilution WB 1:500-2000;IHC-p 1:50-300; ELISA 2000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms -20°C/1 year Observed Band Chromosome . Nucleus, nucleoplasm . Localizes to sites of double-strand breaks (DSBs) following DNA damage. Recruited to DSBs via recognition of RNF168-dependent ubiquitin products Tissue Specificity similarity:Contains 1 RING-type zinc finger., Background Avoid repeated freezing and thawing!	Source	Polyclonal, Rabbit,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms -20°C/1 year Observed Band -20°C/1 year Cell Pathway Chromosome . Nucleus, nucleoplasm . Localizes to sites of double-strand breaks (DSBs) following DNA damage. Recruited to DSBs via recognition of RNF168-dependent ubiquitin products. Tissue Specificity Function Background Avoid repeated freezing and thawing!	Purification	
Purity ≥90% Storage Stability -20°C/1 year Synonyms -20°C/1 year Observed Band -20°C/1 year Cell Pathway Chromosome . Nucleus, nucleoplasm . Localizes to sites of double-strand breaks (DSBs) following DNA damage. Recruited to DSBs via recognition of RNF168-dependent ubiquitin products Tissue Specificity Function Similarity:Contains 1 RING-type zinc finger., Background Avoid repeated freezing and thawing!	Dilution	WB 1:500-2000;IHC-p 1:50-300; ELISA 2000-20000
Storage Stability -20°C/1 year Synonyms -20°C/1 year Observed Band -20°C/1 year Cell Pathway Chromosome . Nucleus, nucleoplasm . Localizes to sites of double-strand breaks (DSBs) following DNA damage. Recruited to DSBs via recognition of RNF168-dependent ubiquitin products Tissue Specificity	Concentration	1 mg/ml
Synonyms Observed Band Cell Pathway Chromosome . Nucleus, nucleoplasm . Localizes to sites of double-strand breaks (DSBs) following DNA damage. Recruited to DSBs via recognition of RNF168-dependent ubiquitin products Tissue Specificity Function Similarity:Contains 1 RING-type zinc finger., Background matters needing Avoid repeated freezing and thawing!	Purity	≥90%
Observed Band Cell Pathway Chromosome . Nucleus, nucleoplasm . Localizes to sites of double-strand breaks (DSBs) following DNA damage. Recruited to DSBs via recognition of RNF168-dependent ubiquitin products Tissue Specificity Function similarity:Contains 1 RING-type zinc finger., Background Matters needing	Storage Stability	-20°C/1 year
Cell Pathway Chromosome . Nucleus, nucleoplasm . Localizes to sites of double-strand breaks (DSBs) following DNA damage. Recruited to DSBs via recognition of RNF168-dependent ubiquitin products Tissue Specificity Function similarity:Contains 1 RING-type zinc finger., Background Avoid repeated freezing and thawing!	Synonyms	
RNF168-dependent ubiquitin products Tissue Specificity Function similarity:Contains 1 RING-type zinc finger., Background matters needing Avoid repeated freezing and thawing!	Observed Band	
Function similarity:Contains 1 RING-type zinc finger., Background matters needing Avoid repeated freezing and thawing!	Cell Pathway	Chromosome . Nucleus, nucleoplasm . Localizes to sites of double-strand breaks (DSBs) following DNA damage. Recruited to DSBs via recognition of RNF168-dependent ubiquitin products
Background matters needing Avoid repeated freezing and thawing!	Tissue Specificity	
matters needing Avoid repeated freezing and thawing!	Function	similarity:Contains 1 RING-type zinc finger.,
matters needing Avoid repeated freezing and thawing! attention	Background	
	matters needing attention	Avoid repeated freezing and thawing!

Nanjing BYabscience technology Co.,Ltd

网址:www.njbybio.com 官方热线:025-5229-8998



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Western blot analysis of lysates from MCF-7 cells, primary antibody was diluted at 1:1000, 4°over night

Immunohistochemical analysis of paraffin-embedded human Breast cancer. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).

Nanjing BYabscience technology Co., Ltd