



HFM1 rabbit pAb

Catalog No BYab-11763 Isotype IgG Reactivity Human;Rat;Mouse; Applications WB Gene Name HFM1 SEC3D1 Protein Name HFM1 Immunogen Synthesized peptide derived from human HFM1 AA range: 943-993 Specificity This antibody detects endogenous levels of HFM1 at Human Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Coll Pathway Tissue Specificity Preferentially expressed in testis and ovary. Function similarity:Belongs to the helicase family, Sk12 subfamily, similarity:Contains 1 helicase C-terminal domain, similarity:Contains 1 SEC63 domain. Background The protein encoded by this gene is thought to be an ATP-dependent DNA helicase and is expressed mainly in germ-line cells. Defects in this gene are a cause of premature ovarian failure 9 (POF9). [provided by RefSeq, Apr 2014], and the provided by RefSeq, Apr 2014].		
Reactivity Human;Rat;Mouse; Applications WB Gene Name HFM1 SEC3D1 Protein Name HFM1 Immunogen Synthesized peptide derived from human HFM1 AA range: 943-993 Specificity This antibody detects endogenous levels of HFM1 at Human Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Tissue Specificity Preferentially expressed in testis and ovary. Function similarity:Belongs to the helicase family, SKI2 subfamily, similarity:Contains 1 helicase ATP-binding domain, similarity-Contains 1 helicase C-terminal domain, similarity:Contains 1 SEC63 domain. Background The protein encoded by this gene is thought to be an ATP-dependent DNA helicase and is expressed mainly in germ-line cells. Defects in this gene are a cause of premature ovarian failure 9 (POF9). [provided by RefSeq, Apr 2014], and the protein of the protein ovarian failure 9 (POF9). [provided by RefSeq, Apr 2014], and the protein ovarian failure 9 (POF9). [provided by RefSeq, Apr 2014], and the protein ovarian failure 9 (POF9). [provided by RefSeq, Apr 2014], and the protein ovarian failure 9 (POF9). [provided by RefSeq, Apr 2014], and the protein ovarian failure 9 (POF9). [provided by RefSeq, Apr 2014], and the protein ovarian failure 9 (POF9). [provided by RefSeq, Apr 2014], and the protein ovarian failure 9 (POF9). [provided by RefSeq, Apr 2014], and the protein ovarian failure 9 (POF9). [provided by RefSeq, Apr 2014], and the protein ovarian failure 9 (POF9). [provided by RefSeq, Apr 2014], and the protein ovarian failure 9 (POF9). [provided by RefSeq, Apr 2014], and the protein ovarian failure 9 (POF9). [provided by RefSeq, Apr 2014], and the protein ovarian failure 9 (POF9). [provided by RefSeq, Apr 2014].	Catalog No	BYab-11763
Applications WB Gene Name HFM1 SEC3D1 Protein Name HFM1 Immunogen Synthesized peptide derived from human HFM1 AA range: 943-993 Specificity This antibody detects endogenous levels of HFM1 at Human Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Tissue Specificity Preferentially expressed in testis and ovary. Function similarity:Belongs to the helicase family. SK12 subfamily.,similarity:Contains 1 helicase ATP-binding domain.,similarity:Contains 1 helicase C-terminal domain.,similarity:Contains 1 SEC63 domain. Background The protein encoded by this gene is thought to be an ATP-dependent DNA helicase and is expressed mainly in germ-line cells. Defects in this gene are a cause of premature ovarian failure 9 (POF9). [provided by RefSeq, Apr 2014], Avoid repeated freezing and thawing!	Isotype	IgG
Gene Name HFM1 SEC3D1 Protein Name HFM1 Immunogen Synthesized peptide derived from human HFM1 AA range: 943-993 Specificity This antibody detects endogenous levels of HFM1 at Human Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Tissue Specificity Preferentially expressed in testis and ovary. Function similarity:Belongs to the helicase family. SKI2 subfamily, similarity:Contains 1 helicase ATP-binding domain., similarity:Contains 1 helicase C-terminal domain., similarity:Contains 1 SEC63 domain., Background The protein encoded by this gene is thought to be an ATP-dependent DNA helicase and is expressed mainly in germ-line cells. Defects in this gene are a cause of premature ovarian failure 9 (POF9). [provided by RefSeq, Apr 2014],	Reactivity	Human;Rat;Mouse;
Protein Name HFM1 Immunogen Synthesized peptide derived from human HFM1 AA range: 943-993 Specificity This antibody detects endogenous levels of HFM1 at Human Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Tissue Specificity Preferentially expressed in testis and ovary. Function similarity:Belongs to the helicase family. SKI2 subfamilysimilarity:Contains 1 helicase ATP-binding domainsimilarity:Contains 1 helicase C-terminal domainsimilarity:Contains 1 SEC63 domain Background The protein encoded by this gene is thought to be an ATP-dependent DNA helicase and is expressed mainly in germ-line cells. Defects in this gene are a cause of premature ovarian failure 9 (POF9). [provided by RefSeq, Apr 2014],	Applications	WB
Immunogen Synthesized peptide derived from human HFM1 AA range: 943-993 Specificity This antibody detects endogenous levels of HFM1 at Human Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Preferentially expressed in testis and ovary. Function similarity:Belongs to the helicase family. SKI2 subfamily.,similarity:Contains 1 helicase ATP-binding domain.,similarity:Contains 1 helicase C-terminal domain.,similarity:Contains 1 SEC63 domain. Background The protein encoded by this gene is thought to be an ATP-dependent DNA helicase and is expressed mainly in germ-line cells. Defects in this gene are a cause of premature ovarian failure 9 (POF9). [provided by RefSeq, Apr 2014], matters needing Avoid repeated freezing and thawing!	Gene Name	HFM1 SEC3D1
Specificity This antibody detects endogenous levels of HFM1 at Human Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Tissue Specificity Preferentially expressed in testis and ovary. Function similarity: Belongs to the helicase family, SKI2 subfamily, similarity: Contains 1 helicase C-terminal domain., similarity: Contains 1 SEC63 domain., Background The protein encoded by this gene is thought to be an ATP-dependent DNA helicase and is expressed mainly in germ-line cells. Defects in this gene are a cause of premature ovarian failure 9 (POF9). [provided by RefSeq, Apr 2014], matters needing Avoid repeated freezing and thawing!	Protein Name	HFM1
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity 290% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Tissue Specificity Preferentially expressed in testis and ovary. Function similarity:Belongs to the helicase family. SKI2 subfamily., similarity:Contains 1 helicase ATP-binding domain., similarity:Contains 1 helicase C-terminal domain., similarity:Contains 1 SEC63 domain., Background The protein encoded by this gene is thought to be an ATP-dependent DNA helicase and is expressed mainly in germ-line cells. Defects in this gene are a cause of premature ovarian failure 9 (POF9). [provided by RefSeq, Apr 2014], matters needing Avoid repeated freezing and thawing!	Immunogen	Synthesized peptide derived from human HFM1 AA range: 943-993
Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Preferentially expressed in testis and ovary. Function similarity:Belongs to the helicase family. SKI2 subfamily.,similarity:Contains 1 helicase ATP-binding domain.,similarity:Contains 1 helicase C-terminal domain.,similarity:Contains 1 SEC63 domain., Background The protein encoded by this gene is thought to be an ATP-dependent DNA helicase and is expressed mainly in germ-line cells. Defects in this gene are a cause of premature ovarian failure 9 (POF9). [provided by RefSeq, Apr 2014], matters needing Avoid repeated freezing and thawing!	Specificity	This antibody detects endogenous levels of HFM1 at Human
Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Tissue Specificity Preferentially expressed in testis and ovary. Function similarity:Belongs to the helicase family. SKI2 subfamily, similarity:Contains 1 helicase ATP-binding domain.,similarity:Contains 1 helicase C-terminal domain.,similarity:Contains 1 SEC63 domain., Background The protein encoded by this gene is thought to be an ATP-dependent DNA helicase and is expressed mainly in germ-line cells. Defects in this gene are a cause of premature ovarian failure 9 (POF9). [provided by RefSeq, Apr 2014], matters needing Avoid repeated freezing and thawing!	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
using specific immunogen. Dilution WB 1: 500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Tissue Specificity Preferentially expressed in testis and ovary. Function similarity:Belongs to the helicase family. SKI2 subfamily.,similarity:Contains 1 helicase ATP-binding domain.,similarity:Contains 1 helicase C-terminal domain.,similarity:Contains 1 SEC63 domain., Background The protein encoded by this gene is thought to be an ATP-dependent DNA helicase and is expressed mainly in germ-line cells. Defects in this gene are a cause of premature ovarian failure 9 (POF9). [provided by RefSeq, Apr 2014], matters needing Avoid repeated freezing and thawing!	Source	Polyclonal, Rabbit,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cell Pathway Tissue Specificity Preferentially expressed in testis and ovary. Function similarity: Belongs to the helicase family. SKI2 subfamily., similarity: Contains 1 helicase ATP-binding domain., similarity: Contains 1 helicase C-terminal domain., similarity: Contains 1 SEC63 domain. Background The protein encoded by this gene is thought to be an ATP-dependent DNA helicase and is expressed mainly in germ-line cells. Defects in this gene are a cause of premature ovarian failure 9 (POF9). [provided by RefSeq, Apr 2014], matters needing Avoid repeated freezing and thawing!	Purification	
Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Tissue Specificity Preferentially expressed in testis and ovary. Function similarity:Belongs to the helicase family. SKI2 subfamily.,similarity:Contains 1 helicase ATP-binding domain.,similarity:Contains 1 helicase C-terminal domain.,similarity:Contains 1 SEC63 domain., Background The protein encoded by this gene is thought to be an ATP-dependent DNA helicase and is expressed mainly in germ-line cells. Defects in this gene are a cause of premature ovarian failure 9 (POF9). [provided by RefSeq, Apr 2014], matters needing Avoid repeated freezing and thawing!	Dilution	WB 1: 500-2000
Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Tissue Specificity Preferentially expressed in testis and ovary. Function similarity:Belongs to the helicase family. SKI2 subfamily.,similarity:Contains 1 helicase ATP-binding domain.,similarity:Contains 1 helicase C-terminal domain.,similarity:Contains 1 SEC63 domain., Background The protein encoded by this gene is thought to be an ATP-dependent DNA helicase and is expressed mainly in germ-line cells. Defects in this gene are a cause of premature ovarian failure 9 (POF9). [provided by RefSeq, Apr 2014], matters needing Avoid repeated freezing and thawing!	Concentration	1 mg/ml
Synonyms Observed Band Cell Pathway Tissue Specificity Preferentially expressed in testis and ovary. Function similarity:Belongs to the helicase family. SKI2 subfamily.,similarity:Contains 1 helicase ATP-binding domain.,similarity:Contains 1 helicase C-terminal domain.,similarity:Contains 1 SEC63 domain., Background The protein encoded by this gene is thought to be an ATP-dependent DNA helicase and is expressed mainly in germ-line cells. Defects in this gene are a cause of premature ovarian failure 9 (POF9). [provided by RefSeq, Apr 2014], matters needing Avoid repeated freezing and thawing!	Purity	≥90%
Observed Band Cell Pathway Tissue Specificity Preferentially expressed in testis and ovary. Function similarity:Belongs to the helicase family. SKI2 subfamily.,similarity:Contains 1 helicase ATP-binding domain.,similarity:Contains 1 helicase C-terminal domain.,similarity:Contains 1 SEC63 domain., Background The protein encoded by this gene is thought to be an ATP-dependent DNA helicase and is expressed mainly in germ-line cells. Defects in this gene are a cause of premature ovarian failure 9 (POF9). [provided by RefSeq, Apr 2014], matters needing Avoid repeated freezing and thawing!	Storage Stability	-20°C/1 year
Tissue Specificity Preferentially expressed in testis and ovary. Function similarity:Belongs to the helicase family. SKI2 subfamily.,similarity:Contains 1 helicase ATP-binding domain.,similarity:Contains 1 helicase C-terminal domain.,similarity:Contains 1 SEC63 domain., Background The protein encoded by this gene is thought to be an ATP-dependent DNA helicase and is expressed mainly in germ-line cells. Defects in this gene are a cause of premature ovarian failure 9 (POF9). [provided by RefSeq, Apr 2014], matters needing Avoid repeated freezing and thawing!	Synonyms	
Tissue Specificity Preferentially expressed in testis and ovary. Similarity:Belongs to the helicase family. SKI2 subfamily.,similarity:Contains 1 helicase ATP-binding domain.,similarity:Contains 1 helicase C-terminal domain.,similarity:Contains 1 SEC63 domain., Background The protein encoded by this gene is thought to be an ATP-dependent DNA helicase and is expressed mainly in germ-line cells. Defects in this gene are a cause of premature ovarian failure 9 (POF9). [provided by RefSeq, Apr 2014], matters needing Avoid repeated freezing and thawing!	Observed Band	
Function similarity:Belongs to the helicase family. SKI2 subfamily.,similarity:Contains 1 helicase ATP-binding domain.,similarity:Contains 1 helicase C-terminal domain.,similarity:Contains 1 SEC63 domain., The protein encoded by this gene is thought to be an ATP-dependent DNA helicase and is expressed mainly in germ-line cells. Defects in this gene are a cause of premature ovarian failure 9 (POF9). [provided by RefSeq, Apr 2014], matters needing Avoid repeated freezing and thawing!	Cell Pathway	
helicase ATP-binding domain., similarity:Contains 1 helicase C-terminal domain., similarity:Contains 1 SEC63 domain., Background The protein encoded by this gene is thought to be an ATP-dependent DNA helicase and is expressed mainly in germ-line cells. Defects in this gene are a cause of premature ovarian failure 9 (POF9). [provided by RefSeq, Apr 2014], matters needing Avoid repeated freezing and thawing!	Tissue Specificity	Preferentially expressed in testis and ovary.
helicase and is expressed mainly in germ-line cells. Defects in this gene are a cause of premature ovarian failure 9 (POF9). [provided by RefSeq, Apr 2014], matters needing Avoid repeated freezing and thawing!	Function	similarity:Belongs to the helicase family. SKI2 subfamily.,similarity:Contains 1 helicase ATP-binding domain.,similarity:Contains 1 helicase C-terminal domain.,similarity:Contains 1 SEC63 domain.,
	Background	The protein encoded by this gene is thought to be an ATP-dependent DNA helicase and is expressed mainly in germ-line cells. Defects in this gene are a
	matters needing attention	Avoid repeated freezing and thawing!

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658




国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from U2OS cells, primary antibody was diluted at 1:1000, 4° over night

Nanjing BYabscience technology Co.,Ltd