



AACS rabbit pAb

	Nanjing BYabscience technology Co.,Ltd
matters needing attention	Avoid repeated freezing and thawing!
Background	
Function	catalytic activity:ATP + acetoacetate + CoA = AMP + diphosphate + acetoacetyl-CoA.,function:Activates acetoacetate to acetoacetyl-CoA. May be involved in utilizing ketone body for the fatty acid-synthesis during adipose tissue development.,similarity:Belongs to the ATP-dependent AMP-binding enzyme family.,tissue specificity:Highly expressed in kidney, heart and brain, but low in liver.,
Tissue Specificity	Highly expressed in kidney, heart and brain, but low in liver.
Cell Pathway	Cytoplasm, cytosol .
Observed Band	
Synonyms	
Storage Stability	-20°C/1 year
Purity	≥90%
Concentration	1 mg/ml
Dilution	WB 1: 500-2000
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Source	Polyclonal, Rabbit,IgG
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Specificity	This antibody detects endogenous levels of AACS at Human/Mouse/Rat
Immunogen	Synthesized peptide derived from human AACS AA range: 193-243
Protein Name	AACS
Gene Name	AACS ACSF1
Applications	WB
Reactivity	Human; Mouse;Rat
Isotype	lgG
Catalog No	BYab-11851

官方热线: 025-5229-8998




国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Nanjing BYabscience technology Co.,Ltd