



SIA4A rabbit pAb

Catalog No	BYab-11853
Isotype	IgG
Reactivity	Human; Mouse
Applications	WB;ELISA;IHC
Gene Name	ST3GAL1 SIAT4 SIAT4A
Protein Name	SIA4A
Immunogen	Synthesized peptide derived from human SIA4A AA range: 189-239
Specificity	This antibody detects endogenous levels of SIA4A at Human/Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:500-2000;IHC-p 1:50-300; ELISA 2000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	
Cell Pathway	Golgi apparatus, Golgi stack membrane ; Single-pass type II membrane protein . Golgi apparatus, trans-Golgi network membrane ; Single-pass type II membrane protein . Secreted. Membrane-bound form in medial and trans cisternae of Golgi (PubMed:9182658). Secreted into the body fluid. .
Tissue Specificity	Expressed in several tissues. Highest expression in lung, liver, skeletal muscle, kidney, pancreas, spleen and placenta.
Function	catalytic activity: CMP-N-acetylneuraminic acid + beta-D-galactosyl-1,3-N-acetyl-alpha-D-galactosaminyl-R = CMP + alpha-N-acetylneuraminyl-2,3-beta-D-galactosyl-1,3-N-acetyl-alpha-D-galactosaminyl-R. function: It may be responsible for the synthesis of the sequence NeuAc-alpha-2,3-Gal-beta-1,3-GalNAc- found on sugar chains O-linked to Thr or Ser and also as a terminal sequence on certain gangliosides. SIAT4A and SIAT4B sialylate the same acceptor substrates but exhibit different Km values. online information: GlycoGene database, online information: ST3Gal I, pathway: Protein modification; protein glycosylation. ,PTM: The soluble form

Nanjing BYabscience technology Co.,Ltd



derives from the membrane form by proteolytic processing.,similarity:Belongs to the glycosyltransferase 29 family.,subcellular location:Membrane-bound form in trans cisternae of Golgi. Secreted into the body fluid.,tissue specificity:Expressed in several tissues. Highest

Background

The protein encoded by this gene is a type II membrane protein that catalyzes the transfer of sialic acid from CMP-sialic acid to galactose-containing substrates. The encoded protein is normally found in the Golgi but can be proteolytically processed to a soluble form. Correct glycosylation of the encoded protein may be critical to its sialyltransferase activity. This protein, which is a member of glycosyltransferase family 29, can use the same acceptor substrates as does sialyltransferase 4B. Two transcript variants encoding the same protein have been found for this gene. Other transcript variants may exist, but have not been fully characterized yet. [provided by RefSeq, Jul 2008],

matters needing attention


Avoid repeated freezing and thawing!

Usage suggestions


This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



Products Images



Western blot analysis of lysates from 293T cells, primary antibody was diluted at 1:1000, 4° over night



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).