



## TIDC1 rabbit pAb

Catalog No	BYab-11913
Isotype	IgG
Reactivity	Human; Mouse;Rat
Applications	WB;ELISA;IHC
Gene Name	TIMMDC1 C3orf1 UNQ247/PRO284
Protein Name	TIDC1
Immunogen	Synthesized peptide derived from human TIDC1 AA range: 74-124
Specificity	This antibody detects endogenous levels of TIDC1 at Human/Mouse/Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:500-2000;IHC-p 1:50-300; ELISA 2000-20000
Concentration	1 mg/ml
Purity	≥90%
Purity Storage Stability	≥90% -20°C/1 year
-	
Storage Stability	
Storage Stability Synonyms	
Storage Stability Synonyms Observed Band	-20°C/1 year
Storage Stability Synonyms Observed Band Cell Pathway	-20°C/1 year Mitochondrion membrane ; Multi-pass membrane protein .
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	-20°C/1 year Mitochondrion membrane ; Multi-pass membrane protein . Generalized expression enhanced in heart and skeletal muscle.
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	-20°C/1 year Mitochondrion membrane ; Multi-pass membrane protein . Generalized expression enhanced in heart and skeletal muscle.
Storage StabilitySynonymsObserved BandCell PathwayTissue SpecificityFunctionBackgroundmatters needing	-20°C/1 year Mitochondrion membrane ; Multi-pass membrane protein . Generalized expression enhanced in heart and skeletal muscle. tissue specificity:Generalized expression enhanced in heart and skeletal muscle.,

## Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658



1

TIDC1

kDa

180 --140 --100 --75 --60 --45 --35 --


25 --

15 --10 --



## **Products Images**

Western blot analysis of lysates from VEC cells, primary antibody was diluted at 1:1000, 4°over night



Immunohistochemical analysis of paraffin-embedded human liver cancer. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).

Nanjing BYabscience technology Co.,Ltd