



NNAT rabbit pAb

Catalog No	BYab-11951
Isotype	IgG
Reactivity	Human; Mouse;Rat
Applications	IHC;IF
Gene Name	NNAT
Protein Name	NNAT
Immunogen	Synthesized peptide derived from human NNAT AA range: 17-67
Specificity	This antibody detects endogenous levels of NNAT at Human/Mouse/Rat
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	IHC-p 1: 50-200. IF 1:50-200
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	
Cell Pathway	cytoplasm,
Tissue Specificity	
Function	developmental stage:Abundant in 18-24 week old fetal brain. Postnatally its expression decline and only minimal levels were present in adulthood.,function:May participate in the maintenance of segment identity in the hindbrain and pituitary development, and maturation or maintenance of the overall structure of the nervous system. May function as a regulatory subunit of ion channels.,similarity:Belongs to the neuronatin family.,
Background	The protein encoded by this gene is a proteolipid that may be involved in the regulation of ion channels during brain development. The encoded protein may also play a role in forming and maintaining the structure of the nervous system. This gene is found within an intron of another gene, bladder cancer associated protein, but on the opposite strand. This gene is imprinted and is expressed only

Nanjing BYabscience technology Co.,Ltd



from the paternal allele. [provided by RefSeq, Apr 2016],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Immunohistochemical analysis of paraffin-embedded human Squamous cell carcinoma of lung. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).