



TTLL6 rabbit pAb

Catalog No	BYab-12029
Isotype	IgG
Reactivity	Human; Mouse
Applications	WB
Gene Name	TTLL6 TTL.6
Protein Name	TTLL6
Immunogen	Synthesized peptide derived from human TTLL6 AA range: 258-308
Specificity	This antibody detects endogenous levels of TTLL6 at Human/Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1: 500-2000
	1 ma/ml
Concentration	1 mg/ml
Concentration Purity	≥90%
	•
Purity	≥90%
Purity Storage Stability	≥90%
Purity Storage Stability Synonyms	≥90%
Purity Storage Stability Synonyms Observed Band	≥90% -20°C/1 year Cytoplasm . Cytoplasm, cytoskeleton . Cytoplasm, cytoskeleton, cilium axoneme . Cytoplasm, cytoskeleton, cilium basal body . CEP41 is required for its transport
Purity Storage Stability Synonyms Observed Band Cell Pathway	≥90% -20°C/1 year Cytoplasm . Cytoplasm, cytoskeleton . Cytoplasm, cytoskeleton, cilium axoneme . Cytoplasm, cytoskeleton, cilium basal body . CEP41 is required for its transport
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	≥90% -20°C/1 year Cytoplasm . Cytoplasm, cytoskeleton . Cytoplasm, cytoskeleton, cilium axoneme . Cytoplasm, cytoskeleton, cilium basal body . CEP41 is required for its transport between the basal body and the cilium axoneme function:Polyglutamylase which preferentially modifies alpha-tubulin. Involved in the side-chain elongation step of the polyglutamylation reaction rather than in the initiation step.,similarity:Belongs to the tubulintyrosine ligase
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	≥90% -20°C/1 year Cytoplasm . Cytoplasm, cytoskeleton . Cytoplasm, cytoskeleton, cilium axoneme . Cytoplasm, cytoskeleton, cilium basal body . CEP41 is required for its transport between the basal body and the cilium axoneme function:Polyglutamylase which preferentially modifies alpha-tubulin. Involved in the side-chain elongation step of the polyglutamylation reaction rather than in the initiation step.,similarity:Belongs to the tubulintyrosine ligase

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658




国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from MDA-MB cells, primary antibody was diluted at 1:1000, 4° over night

Nanjing BYabscience technology Co.,Ltd