



# RIMB1 rabbit pAb

<b>Catalog No</b>	BYab-12237
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human; Mouse;Rat
<b>Applications</b>	WB;IHC
<b>Gene Name</b>	BZRAP1 KIAA0612 RBP1 RIMBP1
<b>Protein Name</b>	RIMB1
<b>Immunogen</b>	Synthesized peptide derived from human RIMB1 AA range: 24-74
<b>Specificity</b>	This antibody detects endogenous levels of RIMB1 at Human/Mouse/Rat
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
<b>Dilution</b>	WB 1:500-2000;IHC-p 1:50-300
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	
<b>Observed Band</b>	
<b>Cell Pathway</b>	Cytoplasm . Mitochondrion . Preferentially expressed in the mitochondria in the presence of TSPO.
<b>Tissue Specificity</b>	Predominantly expressed in brain, pituitary gland and thymus in adults. In adult brain, highest expression found in temporal lobe and the putamen, followed by amygdala, caudate nucleus, cerebral cortex, occipital and frontal lobe. A high expression level is also observed in fetal tissues like brain, heart, kidney and thymus.
<b>Function</b>	caution:PubMed:9915832 demonstrated interaction with BZRP but later PubMed:12435798 demonstrated in the rat ortholog that is not associated with BZRP in the brain.,domain:The SH3 and proline-rich domain is required for the interaction with BZRP and the second SH3 domain mediates binding to a proline-rich motif in RIMS1 and RIMS2.,similarity:Belongs to the RIMBP family.,similarity:Contains 3 fibronectin type-III domains.,similarity:Contains 3 SH3 domains.,subcellular location:Preferentially expressed in the mitochondria in the presence of BZRP.,subunit:Interacts with RIMS1 and RIMS2 (By similarity).

Nanjing BYabscience technology Co.,Ltd



Interacts with BZRP.,tissue specificity:Predominantly expressed in brain, pituitary gland and thymus in adults. In adult brain, highest expression found in temporal lobe and the putamen, followed by amygdala, caudate nucleus, cerebral cortex, occipital and frontal lobe. A high expression level

### Background


### matters needing attention

Avoid repeated freezing and thawing!


### Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images



Western blot analysis of lysates from HeLa cells, primary antibody was diluted at 1:1000, 4° over night



Immunohistochemical analysis of paraffin-embedded human tonsil. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).