



## PHF13 rabbit pAb

Catalog No	BYab-12280
Isotype	IgG
Reactivity	Human; Mouse
Applications	WB;IHC
Gene Name	PHF13
Protein Name	PHF13
Immunogen	Synthesized peptide derived from human PHF13 AA range: 206-256
Specificity	This antibody detects endogenous levels of PHF13 at Human/Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification  Dilution	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.  WB 1:500-2000;IHC-p 1:50-300
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	
Cell Pathway	Nucleus . Nucleus, nucleoplasm . Predominantly bound to chromatin, but a minor proportion is also detected in the nucleoplasm.
Cell Pathway  Tissue Specificity	Nucleus . Nucleus, nucleoplasm . Predominantly bound to chromatin, but a minor proportion is also detected in the nucleoplasm.
•	Nucleus . Nucleus, nucleoplasm . Predominantly bound to chromatin, but a minor proportion is also detected in the nucleoplasm.  similarity:Contains 1 PHD-type zinc finger.,
Tissue Specificity	proportion is also detected in the nucleoplasm.
Tissue Specificity Function	proportion is also detected in the nucleoplasm.
Tissue Specificity Function Background matters needing	proportion is also detected in the nucleoplasm. similarity:Contains 1 PHD-type zinc finger.,


## Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658






## **Products Images**



Western blot analysis of lysates from U2OS cells, primary antibody was diluted at 1:1000, 4° over night



Immunohistochemical analysis of paraffin-embedded human Gastric adenocarcinoma. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).

Nanjing BYabscience technology Co.,Ltd