




IL-17E rabbit pAb

using specific immunogen. Dilution WB 1:1000-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Interleukin-25 (IL-25;Interleukin-17E;IL-17E) Observed Band Cell Pathway Secreted. Tissue Specificity Expressed at low levels in several tissues, including brain, kidney, lung, prostatestis, spinal cord, adrenal gland, and trachea. Function immune system development, leukocyte differentiation, myeloid leukocyte differentiation, defense response, inflammatory response, response to wounding, response to fungus, response to nematode, hemopoiesis, myeloid of the content of the conte		
Reactivity Human;Rat;Mouse; Applications WB; ELISA Gene Name IL25 IL17E UNQ3120/PRO10272 Protein Name IL-17E Immunogen Synthesized peptide derived from human IL-17E AA range: 31-80 Specificity This antibody detects endogenous levels of Human IL-17E Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:1000-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Interleukin-25 (IL-25;Interleukin-17E;IL-17E) Observed Band Cell Pathway Secreted. Tissue Specificity Expressed at low levels in several tissues, including brain, kidney, lung, prostatestis, spinal cord, adrenal gland, and trachea. Immune system development, leukocyte differentiation, myeloid leukocyte differentiation, defense response inflammatory response, response to overlande, hemopolesis, myeloid of differentiation, esinophil differentiation, granulocyte differentiation, hemopolesis or lymphoid organ development, leukocyte differentiation, pranulocyte differentiation, granulocyte differentiation, pranulocyte differentiation, pranulocyte differentiation, granulocyte differentiation, pranulocyte differentiation, granulocyte differentiation, pranulocyte differentiation, granulocyte differentiation, pranulocyte differentiation, pranulocy	Catalog No	BYab-12438
Applications WB; ELISA Gene Name IL-17E Immunogen Synthesized peptide derived from human IL-17E AA range: 31-80 Specificity This antibody detects endogenous levels of Human IL-17E Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:1000-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Interleukin-25 (IL-25;Interleukin-17E;IL-17E) Observed Band Cell Pathway Secreted. Tissue Specificity Expressed at low levels in several tissues, including brain, kidney, lung, prostatestis, spinal cord, adrenal gland, and trachea. Function immune system development, leukocyte differentiation, myeloid leukocyte differentiation, defense response, inflammatory response, response to wounding, response to fungus, response to menadode, hemopoiesis, myeloid or lymphoid organ development, evelopment, granulocyte differentiation, hemopoies or lymphoid organ development, evelopment, granulocyte differentiation, provintiation, provintiation, granulocyte differentiation, provintiation, granulocyte differentiation, provintianting in Tb2-type immune responses, conline information:Interleukin-17 entry, similarity. Belongs to the IL-17 family, tissue specificity. Expressed at low levels in several tissues, including brain, kidney, lung, prostate, testis, spinal cord, adrenal gland, and	Isotype	IgG
Gene Name IL-25 IL17E UNQ3120/PRO10272 Protein Name IL-17E Immunogen Synthesized peptide derived from human IL-17E AA range: 31-80 Specificity This antibody detects endogenous levels of Human IL-17E Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:1000-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Interleukin-25 (IL-25;Interleukin-17E;IL-17E) Observed Band Cell Pathway Cell Pathway Secreted. Tissue Specificity Expressed at low levels in several tissues, including brain, kidney, lung, prostatestis, spinal cord, adrenal gland, and trachea. Function immune system development, leukocyte differentiation, myeloid leukocyte differentiation, defense response, inflammatory response, response to wounding, response to fungus, response to nematode, hemopoiesis, myeloid ordifferentiation, eosinophil differentiation, granulocyte differentiation, hemopoiet or lympoid organ development, eosinophil differentiation, granulocyte differentiation, prionflammatory cy	Reactivity	Human;Rat;Mouse;
IL-17E	Applications	WB; ELISA
Immunogen Synthesized peptide derived from human IL-17E AA range: 31-80 Specificity This antibody detects endogenous levels of Human IL-17E Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:1000-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Interleukin-25 (IL-25;Interleukin-17E;IL-17E) Observed Band Expressed at low levels in several tissues, including brain, kidney, lung, prostatestis, spinal cord, adrenal gland, and trachea. Function Expressed at low levels in several tissues, including brain, kidney, lung, prostated differentiation, development, leukocyte differentiation, myeloid leukocyte differentiation, defense response, inflammatory cepsonse, response to wounding, response to fungus, response to nematode, hemopoiesis, myeloid crifferentiation, essinophil differentiation, granulocyte differentiation, hemopoiet or lymphoid organ development, Background function:Induces activation of NF-kappa-B and stimulates production of Th2-type immune responses, online information:Interleukin-17 entry, similarity:Eelongs to the IL-17 family, tissue specificity:Expressed at low levels in several tissues, includi	Gene Name	IL25 IL17E UNQ3120/PRO10272
Specificity This antibody detects endogenous levels of Human IL-17E Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:1000-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Interleukin-25 (IL-25;Interleukin-17E;IL-17E) Observed Band Cell Pathway Secreted. Tissue Specificity Expressed at low levels in several tissues, including brain, kidney, lung, prostatestis, spinal cord, adrenal gland, and trachea. Function immune system development, leukocyte differentiation, myeloid leukocyte differentiation, defense response, inflammatory response, response to wounding, response to fungus, response to nematode, hemopolesis, myeloid of differentiation, eosinophil differentiation, granulocyte differentiation, hemopolet or lymphoid organ development, Background function:Induces activation of NF-kappa-B and stimulates production of the proinflammatory chemokine IL-8. Proinflammatory cytokine favoring Th2-type immune responses, online information:Interleukin-17 entry, similarity:Belongs to the IL-17 family, tissue specificity:Expressed at low levels in several tissues, including brain, kidney, lung, prostate, testis, spinal cord, adrenal gland, and	Protein Name	IL-17E
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:1000-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Interleukin-25 (IL-25;Interleukin-17E;IL-17E) Observed Band Cell Pathway Secreted. Tissue Specificity Expressed at low levels in several tissues, including brain, kidney, lung, prostatestis, spinal cord, adrenal gland, and trachea. Function immune system development, leukocyte differentiation, myeloid leukocyte differentiation, defense response, inflammatory response, response to wounding, response to fungus, response to nematode, hemopoiesis, myeloid cifferentiation, osoinophil differentiation, granulocyte differentiation, hemopoiet or lymphoid organ development, Background function:Induces activation of NF-kappa-B and stimulates production of the proinflammatory cytokine favoring Th2-type immune responses, online information:Interleukin-17 entry, similarity:Belongs to the IL-17 family, tissue specificity:Expressed at low levels in several tissues, including brain, kidney, lung, prostate, testis, spinal cord, adrenal gland, and	Immunogen	Synthesized peptide derived from human IL-17E AA range: 31-80
Polyclonal, Rabbit, IgG	Specificity	This antibody detects endogenous levels of Human IL-17E
Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:1000-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Interleukin-25 (IL-25;Interleukin-17E;IL-17E) Observed Band Cell Pathway Secreted. Tissue Specificity Expressed at low levels in several tissues, including brain, kidney, lung, prostatestis, spinal cord, adrenal gland, and trachea. Function immune system development, leukocyte differentiation, myeloid leukocyte differentiation, defense response, inflammatory response, response to wounding, response to fungus, response to nematode, hemopoiesis, myeloid or lymphoid organ development, Background Background function:Induces activation of NF-kappa-B and stimulates production of the proinflammatory chemokine IL-8. Proinflammatory cytokine favoring Th2-type immune responses, online information:Interleukin-17 entry, similarity:Belongs to the IL-17 familytissue specificity:Expressed at low levels in several tissues, including brain, kidney, lung, prostate, testis, spinal cord, adrenal gland, and	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
using specific immunogen. Dilution WB 1:1000-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Interleukin-25 (IL-25;Interleukin-17E;IL-17E) Observed Band Cell Pathway Secreted. Tissue Specificity Expressed at low levels in several tissues, including brain, kidney, lung, prostatestis, spinal cord, adrenal gland, and trachea. Function immune system development, leukocyte differentiation, myeloid leukocyte differentiation, defense response to fungus, response to nematode, hemopoiesis, myeloid of differentiation, eosinophil differentiation, granulocyte differentiation, hemopoiet or lymphoid organ development, Background function:Induces activation of NF-kappa-B and stimulates production of the proinflammatory chemokine IL-8. Proinflammatory cytokine favoring Th2-type immune responses.,online information:Interleukin-17 entry, similarity:Belongs to the IL-17 familly_tissue specificity:Expressed at low levels in several tissues, including brain, kidney, lung, prostate, testis, spinal cord, adrenal gland, and	Source	Polyclonal, Rabbit,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Interleukin-25 (IL-25;Interleukin-17E;IL-17E) Observed Band Cell Pathway Secreted. Tissue Specificity Expressed at low levels in several tissues, including brain, kidney, lung, prostatestis, spinal cord, adrenal gland, and trachea. Function immune system development, leukocyte differentiation, myeloid leukocyte differentiation, defense response, inflammatory response, response to wounding, response to fungus, response to nematode, hemopoiesis, myeloid officerentiation, eosinophil differentiation, granulocyte differentiation, hemopoiet or lymphoid organ development, Background function:Induces activation of NF-kappa-B and stimulates production of the proinflammatory chemokine IL-8. Proinflammatory cytokine favoring Th2-type immune responses.,online information:Interleukin-17 entry, similarity:Belongs to the IL-17 familly.,tissue specificity:Expressed at low levels in several tissues, including brain, kidney, lung, prostate, testis, spinal cord, adrenal gland, and	Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Purity ≥90% Storage Stability -20°C/1 year Synonyms Interleukin-25 (IL-25;Interleukin-17E;IL-17E) Observed Band Cell Pathway Secreted. Tissue Specificity Expressed at low levels in several tissues, including brain, kidney, lung, prostatestis, spinal cord, adrenal gland, and trachea. Function immune system development, leukocyte differentiation, myeloid leukocyte differentiation, defense response, inflammatory response, response to wounding, response to fungus, response to nematode, hemopoiesis, myeloid of differentiation, eosinophil differentiation, granulocyte differentiation, hemopoiet or lymphoid organ development, Background function:Induces activation of NF-kappa-B and stimulates production of the proinflammatory chemokine IL-8. Proinflammatory cytokine favoring Th2-type immune responses, online information:Interleukin-17 entry, similarity:Belongs to the IL-17 family, tissue specificity:Expressed at low levels in several tissues, including brain, kidney, lung, prostate, testis, spinal cord, adrenal gland, and	Dilution	WB 1:1000-2000 ELISA 1:5000-20000
Synonyms Interleukin-25 (IL-25;Interleukin-17E;IL-17E) Observed Band Cell Pathway Secreted. Tissue Specificity Expressed at low levels in several tissues, including brain, kidney, lung, prostatestis, spinal cord, adrenal gland, and trachea. Function immune system development, leukocyte differentiation, myeloid leukocyte differentiation, defense response, inflammatory response, response to wounding, response to fungus, response to nematode, hemopoiesis, myeloid of differentiation, eosinophil differentiation, granulocyte differentiation, hemopoiet or lymphoid organ development, Background function:Induces activation of NF-kappa-B and stimulates production of the proinflammatory chemokine IL-8. Proinflammatory cytokine favoring Th2-type immune responses, online information:Interleukin-17 entry, similarity:Belongs to the IL-17 family_tissue specificity:Expressed at low levels in several tissues, including brain, kidney, lung, prostate, testis, spinal cord, adrenal gland, and	Concentration	1 mg/ml
Synonyms Interleukin-25 (IL-25;Interleukin-17E;IL-17E) Observed Band Cell Pathway Secreted. Tissue Specificity Expressed at low levels in several tissues, including brain, kidney, lung, prostatestis, spinal cord, adrenal gland, and trachea. Function immune system development, leukocyte differentiation, myeloid leukocyte differentiation, defense response, inflammatory response, response to wounding, response to fungus, response to nematode, hemopoiesis, myeloid of differentiation, eosinophil differentiation, granulocyte differentiation, hemopoiet or lymphoid organ development, Background function:Induces activation of NF-kappa-B and stimulates production of the proinflammatory chemokine IL-8. Proinflammatory cytokine favoring Th2-type immune responses., online information:Interleukin-17 entry, similarity:Belongs to the IL-17 family., tissue specificity:Expressed at low levels in several tissues, including brain, kidney, lung, prostate, testis, spinal cord, adrenal gland, and		
Cell Pathway Secreted. Tissue Specificity Expressed at low levels in several tissues, including brain, kidney, lung, prostatestis, spinal cord, adrenal gland, and trachea. Function immune system development, leukocyte differentiation, myeloid leukocyte differentiation, defense response, inflammatory response, response to wounding, response to fungus, response to nematode, hemopoiesis, myeloid of differentiation, eosinophil differentiation, granulocyte differentiation, hemopoiet or lymphoid organ development, Background function:Induces activation of NF-kappa-B and stimulates production of the proinflammatory chemokine IL-8. Proinflammatory cytokine favoring Th2-type immune responses., online information:Interleukin-17 entry, similarity:Belongs to the IL-17 family., tissue specificity:Expressed at low levels in several tissues, including brain, kidney, lung, prostate, testis, spinal cord, adrenal gland, and		
Cell Pathway Secreted. Tissue Specificity Expressed at low levels in several tissues, including brain, kidney, lung, prostatestis, spinal cord, adrenal gland, and trachea. Function immune system development, leukocyte differentiation, myeloid leukocyte differentiation, defense response, inflammatory response, response to wounding, response to fungus, response to nematode, hemopoiesis, myeloid of differentiation, eosinophil differentiation, granulocyte differentiation, hemopoiet or lymphoid organ development, Background function:Induces activation of NF-kappa-B and stimulates production of the proinflammatory chemokine IL-8. Proinflammatory cytokine favoring Th2-type immune responses.,online information:Interleukin-17 entry,similarity:Belongs to the IL-17 family.,tissue specificity:Expressed at low levels in several tissues, including brain, kidney, lung, prostate, testis, spinal cord, adrenal gland, and	Purity	≥90%
Tissue Specificity Expressed at low levels in several tissues, including brain, kidney, lung, prostatestis, spinal cord, adrenal gland, and trachea. Function immune system development, leukocyte differentiation, myeloid leukocyte differentiation, defense response, inflammatory response, response to wounding, response to fungus, response to nematode, hemopoiesis, myeloid of differentiation, eosinophil differentiation, granulocyte differentiation, hemopoiet or lymphoid organ development, Background function:Induces activation of NF-kappa-B and stimulates production of the proinflammatory chemokine IL-8. Proinflammatory cytokine favoring Th2-type immune responses.,online information:Interleukin-17 entry,similarity:Belongs to the IL-17 family.,tissue specificity:Expressed at low levels in several tissues, including brain, kidney, lung, prostate, testis, spinal cord, adrenal gland, and	Purity Storage Stability	≥90% -20°C/1 year
Function immune system development, leukocyte differentiation, myeloid leukocyte differentiation, defense response, inflammatory response, response to wounding, response to fungus, response to nematode, hemopoiesis, myeloid of differentiation, eosinophil differentiation, granulocyte differentiation, hemopoiet or lymphoid organ development, Background function:Induces activation of NF-kappa-B and stimulates production of the proinflammatory chemokine IL-8. Proinflammatory cytokine favoring Th2-type immune responses.,online information:Interleukin-17 entry,similarity:Belongs to the IL-17 family.,tissue specificity:Expressed at low levels in several tissues, including brain, kidney, lung, prostate, testis, spinal cord, adrenal gland, and	Purity Storage Stability Synonyms	≥90% -20°C/1 year
differentiation, defense response,inflammatory response, response to wounding, response to fungus, response to nematode, hemopoiesis, myeloid of differentiation, eosinophil differentiation, granulocyte differentiation, hemopoiet or lymphoid organ development, Background function:Induces activation of NF-kappa-B and stimulates production of the proinflammatory chemokine IL-8. Proinflammatory cytokine favoring Th2-type immune responses.,online information:Interleukin-17 entry,similarity:Belongs to the IL-17 family.,tissue specificity:Expressed at low levels in several tissues, including brain, kidney, lung, prostate, testis, spinal cord, adrenal gland, and	Purity Storage Stability Synonyms Observed Band	≥90% -20°C/1 year Interleukin-25 (IL-25;Interleukin-17E;IL-17E)
proinflammatory chemokine IL-8. Proinflammatory cytokine favoring Th2-type immune responses.,online information:Interleukin-17 entry,similarity:Belongs to the IL-17 family.,tissue specificity:Expressed at low levels in several tissues, including brain, kidney, lung, prostate, testis, spinal cord, adrenal gland, and	Purity Storage Stability Synonyms Observed Band Cell Pathway	≥90% -20°C/1 year Interleukin-25 (IL-25;Interleukin-17E;IL-17E) Secreted. Expressed at low levels in several tissues, including brain, kidney, lung, prostate,
	Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	≥90% -20°C/1 year Interleukin-25 (IL-25;Interleukin-17E;IL-17E) Secreted. Expressed at low levels in several tissues, including brain, kidney, lung, prostate, testis, spinal cord, adrenal gland, and trachea. immune system development, leukocyte differentiation, myeloid leukocyte differentiation, defense response,inflammatory response, response to wounding, response to fungus, response to nematode, hemopoiesis, myeloid cell differentiation, eosinophil differentiation, granulocyte differentiation, hemopoietic

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658







attention	Avoid repeated freezing and triawing:
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images