



HP1-β rabbit pAb

Catalog No	BYab-12555
Isotype	IgG
Reactivity	Human; Mouse;Rat
Applications	WB
Gene Name	CBX1 CBX
Protein Name	HP1-β
Immunogen	Synthesized peptide derived from human HP1-β
Specificity	This antibody detects endogenous levels of HP1-β at Human, Mouse,Rat
Formulation	Liquid in PBS containing 50% glycerol, and 0.43% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	Chromobox protein homolog 1 (HP1Hsbeta) (Heterochromatin protein 1 homolog beta) (HP1 beta) (Heterochromatin protein p25) (M31) (Modifier 1 protein) (p25beta)
Observed Band	
Cell Pathway	Nucleus . Unassociated with chromosomes during mitosis.
Tissue Specificity	Expressed in all adult and embryonic tissues.
Function	function:Component of heterochromatin. Recognizes and binds histone H3 tails methylated at 'Lys-9', leading to epigenetic repression. Interaction with lamin B receptor (LBR) can contribute to the association of the heterochromatin with the inner nuclear membrane.,online information:Heterochromatin protein 1 entry,PTM:Not phosphorylated.,similarity:Contains 2 chromo domains.,subcellular location:Unassociated with chromosomes during mitosis.,subunit:Homodimer. Interacts directly with CHAF1A, EMSY, LBR, TIF1/TIF1A and TRIM28/TIF1B PXVXL motif via the chromoshadow domain. Interacts directly with histone H3 methylated at 'Lys-9' via the chromo domain. Interacts with SUV39H1 and SETDB1, SUV420H1 and SUV420H2. Interacts with PRDM6.,tissue specificity:In

Nanjing BYabscience technology Co.,Ltd



all adult and embryonic tissues.,

Background

This gene encodes a highly conserved nonhistone protein, which is a member of the heterochromatin protein family. The protein is enriched in the heterochromatin and associated with centromeres. The protein has a single N-terminal chromodomain which can bind to histone proteins via methylated lysine residues, and a C-terminal chromo shadow-domain (CSD) which is responsible for the homodimerization and interaction with a number of chromatin-associated nonhistone proteins. The protein may play an important role in the epigenetic control of chromatin structure and gene expression. Several related pseudogenes are located on chromosomes 1, 3, and X. Multiple alternatively spliced variants, encoding the same protein, have been identified. [provided by RefSeq, Jul 2008],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images