



VAMP4 rabbit pAb

Catalog No BYab-12557 Isotype IgG Reactivity Human; Mouse;Rat Applications WB Gene Name VAMP4 Protein Name VAMP4 Immunogen Synthesized peptide derived from human VAMP4 Specificity This antibody detects endogenous levels of VAMP4 at Human, Mouse,Rat Formulation Liquid in PBS containing 50% glycerol, and 0.47% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Vesicle-associated membrane protein 4 (VAMP-4) Observed Band 17kD Cell Pathway Golgi apparatus, trans-Golgi network membrane; Single-pass type IV membran profein. Associated with trans Golgi network (TGN) and newly formed immature secretory granules (ISG.) Not found on the mature secretory organelles. Tissue Specificity B-cell,Bone marrow,Epithelium,Liver,Urinary bladder, Function function:Involved in the pathway that functio		
Reactivity Human; Mouse;Rat Applications WB Gene Name VAMP4 Protein Name VAMP4 Immunogen Synthesized peptide derived from human VAMP4 Specificity This antibody detects endogenous levels of VAMP4 at Human, Mouse,Rat Formulation Liquid in PBS containing 50% glycerol, and 0.47% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Vesicle-associated membrane protein 4 (VAMP-4) Observed Band 17kD Cell Pathway Golgi apparatus, trans-Golgi network membrane ; Single-pass type IV membrang protein . Associated with trans Golgi network (TGN) and newly formed immature secretory granules (ISG). Not found on the mature secretory organelles. Tissue Specificity B-cell,Bone marrow,Epithelium,Liver,Urinary bladder, function:Involved in the pathway that functions to remove an inhibitor (probably synaptotagmin-4) of calcium-triggered exocytosis during the maturation of secretory granules. May be a marker for this sorting pathway that is critical for remodeling the secretory granules. May be a marker for this sorting pathway that is critical for remodeling the secretory granules. May be a marker for this sorting pathway that is critical for remodeling the secretory granules. May be a marker for this sorting pathway that is critical for remodeling the secretory granules. May be a marker for this sorting pathway that is critical for remodeling the secretory granules. May be a marker for this sorting pathway that is critical for remodeling the secretory organelles. Subunit:Identified in a complex containing STX6, STX13,	Catalog No	BYab-12557
Applications WB Gene Name VAMP4 Protein Name VAMP4 Immunogen Synthesized peptide derived from human VAMP4 Specificity This antibody detects endogenous levels of VAMP4 at Human, Mouse,Rat Formulation Liquid in PBS containing 50% glycerol, and 0.47% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Vesicle-associated membrane protein 4 (VAMP-4) Observed Band 17kD Cell Pathway Golgi apparatus, trans-Golgi network membrane: Single-pass type IV membrar protein. Associated with trans Golgi network (TGN) and newly formed immature secretory granules (ISG). Not found on the mature secretory organelles. Tissue Specificity B-cell,Bone marrow,Epithelium,Liver,Urinary bladder, function:Involved in the pathway that functions to remove an inhibitor (probably synaptotagmin-4) of calcium-triggered exocytosis during the maturation of secretory granules. May be a marker for this sorting pathway that is critical for remodeling the secretory response of granule, similarity:Belongs to the synaptobrevin family, similarity:Contains 1 v-SNARE coiled-coil homology domain, subcellular location:Associated with trans Colgi network (TGN) and newly formed immature secretory granules (ISG). Not found on the mature secretory organelles, subunit-lidentified in a complex containing STX6, STX13,	Isotype	IgG
Gene Name VAMP4 Protein Name VAMP4 Immunogen Synthesized peptide derived from human VAMP4 Specificity This antibody detects endogenous levels of VAMP4 at Human, Mouse,Rat Formulation Liquid in PBS containing 50% glycerol, and 0.47% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Vesicle-associated membrane protein 4 (VAMP-4) Observed Band 17kD Cell Pathway Golgi apparatus, trans-Golgi network membrane; Single-pass type IV membrar protein . Associated with trans Golgi network (TGN) and newly formed immature secretory granules (ISG). Not found on the mature secretory organelles. Tissue Specificity B-cell,Bone marrow,Epithelium,Liver,Urinary bladder, function:Involved in the pathway that functions to remove an inhibitor (probably synaptotagmin-4) of calcium-triggered exceytosis during the maturation of secretory granules. May be a marker for this sorting pathway that is critical for remodeling the secretory granules should resource the more of the sorting pathway that is critical for remodeling the secretory granules (ISG). Not found on the motowork (TGN) and newly formed immature secretory granules (ISG). Not found on the mature in the mature secretory organelles. Subunit-Identified in a complex containing STX6, STX13,	Reactivity	Human; Mouse;Rat
Protein Name VAMP4 Immunogen Synthesized peptide derived from human VAMP4 Specificity This antibody detects endogenous levels of VAMP4 at Human, Mouse,Rat Formulation Liquid in PBS containing 50% glycerol, and 0.47% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Vesicle-associated membrane protein 4 (VAMP-4) Observed Band 17kD Cell Pathway Golgi apparatus, trans-Golgi network membrane: Single-pass type IV membran protein . Associated with trans Golgi network (TGN) and newly formed immature secretory granules (ISG). Not found on the mature secretory organelles. Tissue Specificity B-cell,Bone marrow,Epithelium,Liver,Urinary bladder, function:Involved in the pathway that functions to remove an inhibitor (probably synaptotagmin-4) of calcium-triggered exocytosis during the maturation of secretory granules. May be a marker for this sorting pathway that is critical for remodeling the secretory response of granule. similarity:Belongs to the synaptotagmin family. similarity:Contains 1 v-SNARE coiled-coil homology domain. subcellular location:Associated with trans Golgi network (TGN) and newly formed immature secretory granules (ISG). Not found on the mature secretory organelles. Subunitied on a complex containing STX6, STX13,	Applications	WB
Immunogen Synthesized peptide derived from human VAMP4 Specificity This antibody detects endogenous levels of VAMP4 at Human, Mouse,Rat Formulation Liquid in PBS containing 50% glycerol, and 0.47% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Vesicle-associated membrane protein 4 (VAMP-4) Observed Band 17kD Cell Pathway Golgi apparatus, trans-Golgi network membrane : Single-pass type IV membrar protein . Associated with trans Golgi network (TGN) and newly formed immature secretory granules (ISG). Not found on the mature secretory organelles. Tissue Specificity B-cell,Bone marrow,Epithelium,Liver,Urinary bladder, Function function:Involved in the pathway that functions to remove an inhibitor (probably synaptotagmin-4) of calcium-triggered exocytosis during the maturation of secretory granules. May be a marker for this sorting pathway that is critical for remodeling the secretory response of granule, similarity:Belongs to the synaptobrevin family, similarity. Contains 1 v-SNARE collect-coll homology domain, subcellular location:Associated with trans Golgi network (TGN) and newly formed i	Gene Name	VAMP4
Formulation Liquid in PBS containing 50% glycerol, and 0.47% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Vesicle-associated membrane protein 4 (VAMP-4) Observed Band 17kD Cell Pathway Golgi apparatus, trans-Golgi network membrane; Single-pass type IV membran protein. Associated with trans Golgi network (TGN) and newly formed immature secretory granules (ISG). Not found on the mature secretory organelles. Tissue Specificity Function function:Involved in the pathway that functions to remove an inhibitor (probably synaptotagmin-4) of calcium-triggered exocyctosis during the maturation of secretory granules. May be a marker for this sorting pathway that is critical for remodeling the secretory response of granule, similarity:Belongs to the synaptobrevin family, similarity:Contains 1 v-SNARE coiled-coil homology domain.,subcellular location:Associated with trans Golgi network (TGN) and newly formed immature secretory granules (ISG). Not found on the matures of granule, similarity:Belongs to the synaptobrevin family, similarity:Contains 1 v-SNARE coiled-coil homology domain, subcellular location:Associated with trans Golgi network (TGN) and newly formed immature secretory granules (ISG). Not found on the mature secretory organelles, subunit itidentified in a complex containing STX6, STX13,	Protein Name	VAMP4
Formulation Liquid in PBS containing 50% glycerol, and 0.47% sodium azide. Source Polyclonal, Rabbit, IgG The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Vesicle-associated membrane protein 4 (VAMP-4) Observed Band 17kD Cell Pathway Golgi apparatus, trans-Golgi network membrane; Single-pass type IV membran protein. Associated with trans Golgi network (TGN) and newly formed immature secretory granules (ISG). Not found on the mature secretory organelles. Tissue Specificity Function function:Involved in the pathway that functions to remove an inhibitor (probably synaptotagmin-4) of calcium-triggered exocytosis during the maturation of secretory granules. May be a marker for this sorting pathway that is critical for remodeling the secretory response of granule. similarity:Belongs to the synaptobrevin family, similarity:Contains 1 v-SNARE coiled-coil homology domain., subcellular location:Associated with trans Golgi network (TGN) and newly formed immature secretory granules (ISG). Not found on the mature secretory organelles. subunit:Identlifed in a complex containing STX6, STX13,	Immunogen	Synthesized peptide derived from human VAMP4
Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Vesicle-associated membrane protein 4 (VAMP-4) Observed Band 17kD Cell Pathway Golgi apparatus, trans-Golgi network membrane; Single-pass type IV membrang protein. Associated with trans Golgi network (TGN) and newly formed immature secretory granules (ISG). Not found on the mature secretory organelles. Tissue Specificity B-cell,Bone marrow,Epithelium,Liver,Urinary bladder, Function function:Involved in the pathway that functions to remove an inhibitor (probably synaptotagmin-4) of calcium-triggered exocytosis during the maturation of secretory granules. May be a marker for this sorting pathway that is critical for remodeling the secretory response of granule, similarity:Belongs to the synaptobrevin family. similarity:Contains 1 v-SNARE coiled-coil homology domain, subcellular location:Associated with trans Golgi network (TGN) and newly formed immature secretory granules (ISG). Not found on the mature secretory granu	Specificity	This antibody detects endogenous levels of VAMP4 at Human, Mouse,Rat
Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Vesicle-associated membrane protein 4 (VAMP-4) Observed Band 17kD Cell Pathway Golgi apparatus, trans-Golgi network membrane; Single-pass type IV membran protein. Associated with trans Golgi network (TGN) and newly formed immature secretory granules (ISG). Not found on the mature secretory organelles. Tissue Specificity B-cell,Bone marrow,Epithelium,Liver,Urinary bladder, function:Involved in the pathway that functions to remove an inhibitor (probably synaptotagmin-4) of calcium-triggered exocytosis during the maturation of secretory granules. May be a marker for this sorting pathway that is critical for remodelling the secretory response of granule., similarity:Belongs to the synaptobrevin family.,similarity:Contains 1 v-SNARE coiled-coil homology domain.,subcellular location:Associated with trans Golgi network (TGN) and newly formed immature secretory granules (ISG). Not found on the mature secretory organelles.,subunit:Identified in a complex containing STX6, STX13,	Formulation	Liquid in PBS containing 50% glycerol, and 0.47% sodium azide.
using specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Vesicle-associated membrane protein 4 (VAMP-4) Observed Band 17kD Cell Pathway Golgi apparatus, trans-Golgi network membrane; Single-pass type IV membrar protein . Associated with trans Golgi network (TGN) and newly formed immature secretory granules (ISG). Not found on the mature secretory organelles. Tissue Specificity B-cell,Bone marrow,Epithelium,Liver,Urinary bladder, Function function:Involved in the pathway that functions to remove an inhibitor (probably synaptotagmin-4) of calcium-triggered exocytosis during the maturation of secretory granules. May be a marker for this sorting pathway that is critical for remodeling the secretory response of granule, similarity;Belongs to the synaptobrevin family, similarity:Contains 1 v-SNARE coiled-coil homology domain, subcellular location:Associated with trans Golgi network (TGN) and newly formed immature secretory granules (ISG). Not found on the mature secretory organelless, subunit:Identified in a complex containing STX6, STX13,	Source	Polyclonal, Rabbit,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Vesicle-associated membrane protein 4 (VAMP-4) Observed Band 17kD Cell Pathway Golgi apparatus, trans-Golgi network membrane; Single-pass type IV membran protein. Associated with trans Golgi network (TGN) and newly formed immature secretory granules (ISG). Not found on the mature secretory organelles. Tissue Specificity B-cell,Bone marrow,Epithelium,Liver,Urinary bladder, Function function:Involved in the pathway that functions to remove an inhibitor (probably synaptotagmin-4) of calcium-triggered exocytosis during the maturation of secretory granules. May be a marker for this sorting pathway that is critical for remodeling the secretory response of granule.,similarity:Belongs to the synaptobrevin family.,similarity:Contains 1 v-SNARE coiled-coil homology domain.,subcellular location:Associated with trans Golgi network (TGN) and newly formed immature secretory granules (ISG). Not found on the mature secretory organelles.,subunit:Identified in a complex containing STX6, STX13,	Purification	
Purity ≥90% Storage Stability -20°C/1 year Synonyms Vesicle-associated membrane protein 4 (VAMP-4) Observed Band 17kD Cell Pathway Golgi apparatus, trans-Golgi network membrane; Single-pass type IV membrar protein. Associated with trans Golgi network (TGN) and newly formed immature secretory granules (ISG). Not found on the mature secretory organelles. Tissue Specificity B-cell,Bone marrow,Epithelium,Liver,Urinary bladder, Function function:Involved in the pathway that functions to remove an inhibitor (probably synaptotagmin-4) of calcium-triggered exocytosis during the maturation of secretory granules. May be a marker for this sorting pathway that is critical for remodeling the secretory response of granule.,similarity:Belongs to the synaptobrevin family.,similarity:Contains 1 v-SNARE coiled-coil homology domain.,subcellular location:Associated with trans Golgi network (TGN) and newly formed immature secretory granules (ISG). Not found on the mature secretory organelles.,subunit:Identified in a complex containing STX6, STX13,	Dilution	WB 1:500-2000
Storage Stability -20°C/1 year Vesicle-associated membrane protein 4 (VAMP-4) Observed Band 17kD Golgi apparatus, trans-Golgi network membrane; Single-pass type IV membrare protein. Associated with trans Golgi network (TGN) and newly formed immature secretory granules (ISG). Not found on the mature secretory organelles. Tissue Specificity B-cell,Bone marrow,Epithelium,Liver,Urinary bladder, function:Involved in the pathway that functions to remove an inhibitor (probably synaptotagmin-4) of calcium-triggered exocytosis during the maturation of secretory granules. May be a marker for this sorting pathway that is critical for remodeling the secretory response of granule.,similarity:Belongs to the synaptobrevin family.,similarity:Contains 1 v-SNARE coiled-coil homology domain.,subcellular location:Associated with trans Golgi network (TGN) and newly formed immature secretory granules (ISG). Not found on the mature secretory organelles.,subunit:Identified in a complex containing STX6, STX13,	Concentration	1 mg/ml
Synonyms Vesicle-associated membrane protein 4 (VAMP-4) Observed Band 17kD Cell Pathway Golgi apparatus, trans-Golgi network membrane; Single-pass type IV membrare protein. Associated with trans Golgi network (TGN) and newly formed immature secretory granules (ISG). Not found on the mature secretory organelles. B-cell,Bone marrow,Epithelium,Liver,Urinary bladder, function function:Involved in the pathway that functions to remove an inhibitor (probably synaptotagmin-4) of calcium-triggered exocytosis during the maturation of secretory granules. May be a marker for this sorting pathway that is critical for remodeling the secretory response of granulesimilarity:Belongs to the synaptobrevin familysimilarity:Contains 1 v-SNARE coiled-coil homology domainsubcellular location:Associated with trans Golgi network (TGN) and newly formed immature secretory granules (ISG). Not found on the mature secretory organellessubunit:Identified in a complex containing STX6, STX13,	Purity	≥90%
Observed Band Cell Pathway Golgi apparatus, trans-Golgi network membrane; Single-pass type IV membrare protein. Associated with trans Golgi network (TGN) and newly formed immature secretory granules (ISG). Not found on the mature secretory organelles. B-cell,Bone marrow,Epithelium,Liver,Urinary bladder, function:Involved in the pathway that functions to remove an inhibitor (probably synaptotagmin-4) of calcium-triggered exocytosis during the maturation of secretory granules. May be a marker for this sorting pathway that is critical for remodeling the secretory response of granule.,similarity:Belongs to the synaptobrevin family.,similarity:Contains 1 v-SNARE coiled-coil homology domain.,subcellular location:Associated with trans Golgi network (TGN) and newly formed immature secretory granules (ISG). Not found on the mature secretory organelles.,subunit:Identified in a complex containing STX6, STX13,	Storage Stability	-20°C/1 year
Cell Pathway Golgi apparatus, trans-Golgi network membrane; Single-pass type IV membrare protein. Associated with trans Golgi network (TGN) and newly formed immature secretory granules (ISG). Not found on the mature secretory organelles. B-cell,Bone marrow,Epithelium,Liver,Urinary bladder, function:Involved in the pathway that functions to remove an inhibitor (probably synaptotagmin-4) of calcium-triggered exocytosis during the maturation of secretory granules. May be a marker for this sorting pathway that is critical for remodeling the secretory response of granule.,similarity:Belongs to the synaptobrevin family.,similarity:Contains 1 v-SNARE coiled-coil homology domain.,subcellular location:Associated with trans Golgi network (TGN) and newly formed immature secretory granules (ISG). Not found on the mature secretory organelles.,subunit:Identified in a complex containing STX6, STX13,	Synonyms	Vesicle-associated membrane protein 4 (VAMP-4)
Tissue Specificity B-cell,Bone marrow,Epithelium,Liver,Urinary bladder, function function:Involved in the pathway that functions to remove an inhibitor (probably synaptotagmin-4) of calcium-triggered exocytosis during the maturation of secretory granules. May be a marker for this sorting pathway that is critical for remodeling the secretory response of granule.,similarity:Belongs to the synaptobrevin family.,similarity:Contains 1 v-SNARE coiled-coil homology domain.,subcellular location:Associated with trans Golgi network (TGN) and newly formed immature secretory granules (ISG). Not found on the mature secretory organelles.,subunit:Identified in a complex containing STX6, STX13,	Observed Band	17kD
function:Involved in the pathway that functions to remove an inhibitor (probably synaptotagmin-4) of calcium-triggered exocytosis during the maturation of secretory granules. May be a marker for this sorting pathway that is critical for remodeling the secretory response of granule.,similarity:Belongs to the synaptobrevin family.,similarity:Contains 1 v-SNARE coiled-coil homology domain.,subcellular location:Associated with trans Golgi network (TGN) and newly formed immature secretory granules (ISG). Not found on the mature secretory organelles.,subunit:Identified in a complex containing STX6, STX13,	Cell Pathway	Golgi apparatus, trans-Golgi network membrane; Single-pass type IV membrane protein. Associated with trans Golgi network (TGN) and newly formed immature secretory granules (ISG). Not found on the mature secretory organelles.
synaptotagmin-4) of calcium-triggered exocytosis during the maturation of secretory granules. May be a marker for this sorting pathway that is critical for remodeling the secretory response of granule.,similarity:Belongs to the synaptobrevin family.,similarity:Contains 1 v-SNARE coiled-coil homology domain.,subcellular location:Associated with trans Golgi network (TGN) and newly formed immature secretory granules (ISG). Not found on the mature secretory organelles.,subunit:Identified in a complex containing STX6, STX13,	Tissue Specificity	B-cell,Bone marrow,Epithelium,Liver,Urinary bladder,
	Function	synaptotagmin-4) of calcium-triggered exocytosis during the maturation of secretory granules. May be a marker for this sorting pathway that is critical for remodeling the secretory response of granule.,similarity:Belongs to the synaptobrevin family.,similarity:Contains 1 v-SNARE coiled-coil homology domain.,subcellular location:Associated with trans Golgi network (TGN) and newly formed immature secretory granules (ISG). Not found on the mature secretory organelles.,subunit:Identified in a complex containing STX6, STX13,

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658







Background	Synaptobrevins/VAMPs, syntaxins, and the 25-kD synaptosomal-associated protein SNAP25 are the main components of a protein complex involved in the docking and/or fusion of synaptic vesicles with the presynaptic membrane. The protein encoded by this gene is a member of the vesicle-associated membrane protein (VAMP)/synaptobrevin family. This protein may play a role in trans-Golgi network-to-endosome transport. [provided by RefSeq, Jul 2008],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658