



# MAGI-2 Polyclonal Antibody

<b>Catalog No</b>	BYab-12744
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Mouse;Rat
<b>Applications</b>	WB;IHC;IF;ELISA
<b>Gene Name</b>	MAGI2
<b>Protein Name</b>	Membrane-associated guanylate kinase WW and PDZ domain-containing protein 2
<b>Immunogen</b>	The antiserum was produced against synthesized peptide derived from human MAGI2. AA range:221-270
<b>Specificity</b>	MAGI-2 Polyclonal Antibody detects endogenous levels of MAGI-2 protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB: 1/500 - 1/2000. IHC: 1/100 - 1/300. ELISA: 1/20000.. IF 1:50-200
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	MAGI2; ACVRINP1; AIP1; KIAA0705; Membrane-associated guanylate kinase; WW and PDZ domain-containing protein 2; Atrophin-1-interacting protein 1; AIP-1; Atrophin-1-interacting protein A; Membrane-associated guanylate kinase inverted 2; MAGI-
<b>Observed Band</b>	156kD
<b>Cell Pathway</b>	Cytoplasm . Late endosome . Cell junction, synapse, synaptosome . Cell membrane ; Peripheral membrane protein . Localized diffusely in the cytoplasm before nerve growth factor (NGF) stimulation. Recruited to late endosomes after NGF stimulation. Membrane-associated in synaptosomes (By similarity). .
<b>Tissue Specificity</b>	Specifically expressed in brain.
<b>Function</b>	function:Seems to act as scaffold molecule at synaptic junctions by assembling neurotransmitter receptors and cell adhesion proteins. May play a role in regulating activin-mediated signaling in neuronal cells. Enhances the ability of PTEN to suppress AKT1 activation.,similarity:Belongs to the MAGUK family.,similarity:Contains 1 guanylate kinase-like domain.,similarity:Contains 2

Nanjing BYabscience technology Co.,Ltd



WW domains.,similarity:Contains 6 PDZ (DHR) domains.,subcellular location:Membrane-associated in synaptosomes.,subunit:Interacts via its WW domains with DRPLA. Interacts via its second PDZ domain with PTEN unphosphorylated C-terminus; this interaction diminishes the degradation rate of PTEN (By similarity). Interacts through its guanylate kinase domain with DLGAP1 (By similarity). Interacts through the PDZ domains with GRIN2A, GRID2 and NLGN1 (By similarity). Interacts with CTNND2, CTNNB1, MAGUIN-1, ACVR2A, SMAD2

#### Background

The protein encoded by this gene interacts with atrophin-1. Atrophin-1 contains a polyglutamine repeat, expansion of which is responsible for dentatorubral and pallidoluysian atrophy. This encoded protein is characterized by two WW domains, a guanylate kinase-like domain, and multiple PDZ domains. It has structural similarity to the membrane-associated guanylate kinase homologue (MAGUK) family. [provided by RefSeq, Jul 2008],

#### matters needing attention

Avoid repeated freezing and thawing!


#### Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.


Nanjing BYabscience technology Co.,Ltd




## Products Images




Ma, Mengtao, et al. "miR-487a promotes TGF-β 1-induced EMT, the migration and invasion of breast cancer cells by directly targeting MAGI2." International journal of biological sciences 12.4 (2016): 397.



Immunohistochemical analysis of paraffin-embedded Human breast cancer. Antibody was diluted at 1:100(4° overnight). High-pressure and temperature Tris-EDTA,pH8.0 was used for antigen retrieval. Negative contrl (right) obtained from antibody was pre-absorbed by immunogen peptide.



Western blot analysis of lysates from A549 cells, using MAGI2 Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from 293 cells using MAGI2 antibody.