



# IL-9R (phospho Ser519) Polyclonal Antibody

<b>Catalog No</b>	BYab-13057
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Rat;Mouse;
<b>Applications</b>	WB;IHC;IF;ELISA
<b>Gene Name</b>	IL9R
<b>Protein Name</b>	Interleukin-9 receptor
<b>Immunogen</b>	The antiserum was produced against synthesized peptide derived from human IL-9R around the phosphorylation site of Ser519. AA range:472-521
<b>Specificity</b>	Phospho-IL-9R (S519) Polyclonal Antibody detects endogenous levels of IL-9R protein only when phosphorylated at S519.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications.
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	IL9R; Interleukin-9 receptor; IL-9 receptor; IL-9R; CD antigen CD129
<b>Observed Band</b>	57kD
<b>Cell Pathway</b>	Cell membrane; Single-pass type I membrane protein. Secreted.
<b>Tissue Specificity</b>	Melanoma,
<b>Function</b>	domain:The box 1 motif is required for JAK interaction and/or activation.,domain:The WSXWS motif appears to be necessary for proper protein folding and thereby efficient intracellular transport and cell-surface receptor binding.,function:This is a receptor for interleukin-9.,miscellaneous:The gene encoding for this protein is located in the pseudoautosomal region 2 (PAR2) of X and Y chromosomes.,similarity:Belongs to the type I cytokine receptor family. Type 4 subfamily.,similarity:Contains 1 fibronectin type-III domain.,
<b>Background</b>	The protein encoded by this gene is a cytokine receptor that specifically mediates the biological effects of interleukin 9 (IL9). The functional IL9 receptor

Nanjing BYabscience technology Co.,Ltd



complex requires this protein as well as the interleukin 2 receptor, gamma (IL2RG), a common gamma subunit shared by the receptors of many different cytokines. The ligand binding of this receptor leads to the activation of various JAK kinases and STAT proteins, which connect to different biologic responses. This gene is located at the pseudoautosomal regions of X and Y chromosomes. Genetic studies suggested an association of this gene with the development of asthma. Multiple pseudogenes on chromosome 9, 10, 16, and 18 have been described. Alternatively spliced transcript variants have been found for this gene. [provided by RefSeq, Jul 2008],

**matters needing attention**


Avoid repeated freezing and thawing!

**Usage suggestions**


This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.




## Products Images




Enzyme-Linked Immunosorbent Assay (Phospho-ELISA) for Immunogen Phosphopeptide (Phospho-left) and Non-Phosphopeptide (Phospho-right), using IL-9R (Phospho-Ser519) Antibody



Immunofluorescence analysis of HUVEC cells, using IL-9R (Phospho-Ser519) Antibody. The picture on the right is blocked with the phospho peptide.



Immunohistochemistry analysis of paraffin-embedded human lymph node, using IL-9R (Phospho-Ser519) Antibody. The picture on the right is blocked with the phospho peptide.



Western blot analysis of lysates from HUVEC cells, using IL-9R (Phospho-Ser519) Antibody. The lane on the right is blocked with the phospho peptide.

Nanjing BYabscience technology Co.,Ltd