




GPR40 Polyclonal Antibody

| Catalog No | BYab-13337 |
|--------------------|---|
| Isotype | IgG |
| Reactivity | Human;Monkey |
| Applications | WB;IF;ELISA |
| Gene Name | FFAR1 |
| Protein Name | Free fatty acid receptor 1 |
| Immunogen | The antiserum was produced against synthesized peptide derived from human FFAR1. AA range:185-234 |
| Specificity | GPR40 Polyclonal Antibody detects endogenous levels of GPR40 protein. |
| Formulation | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |
| Source | Polyclonal, Rabbit,IgG |
| Purification | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Dilution | Western Blot: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/10000. Not yet tested in other applications. |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/1 year |
| Synonyms | FFAR1; GPR40; Free fatty acid receptor 1; G-protein coupled receptor 40 |
| Observed Band | 26kD |
| Cell Pathway | Cell membrane ; Multi-pass membrane protein . |
| Tissue Specificity | Detected in brain and pancreas. Detected in pancreatic beta cells. |
| Function | function:Receptor for medium and long chain saturated and unsaturated fatty acids. Binding of the ligand increase intracellular calcium concentration and amplify glucose-stimulated insulin secretion. The activity of this receptor is mediated by G-proteins that activate phospholipase C. Seems to act through a G(q) and G(i)-mediated pathway.,similarity:Belongs to the G-protein coupled receptor 1 family.,tissue specificity:Expressed abundantly in pancreatic beta cells., |
| Background | This gene encodes a member of the GP40 family of G protein-coupled receptors that are clustered together on chromosome 19. The encoded protein is a receptor for medium and long chain free fatty acids and may be involved in the metabolic |

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658








| | regulation of insulin secretion. Polymorphisms in this gene may be associated with type 2 diabetes. [provided by RefSeq, Apr 2009], |
|---------------------------|---|
| matters needing attention | Avoid repeated freezing and thawing! |
| Usage suggestions | This product can be used in immunological reaction related experiments. For more information, please consult technical personnel. |
| | |







Products Images




Western Blot analysis of COS7 cells using GPR40 Polyclonal Antibody diluted at 1:500



Western Blot analysis of COS-7 cells using GPR40 Polyclonal Antibody diluted at 1:500



Western blot analysis of Mouse-kidney lysis using GPR40 antibody. Antibody was diluted at 1:500



Immunofluorescence analysis of LOVO cells, using FFAR1 Antibody. The picture on the right is blocked with the synthesized peptide.

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



