



## IL-17Rα Polyclonal Antibody

| Catalog No         | BYab-13374   |
|--------------------|--|
| Isotype            | IgG  |
| Reactivity         | Human;Mouse  |
| Applications       | WB;ELISA   |
| Gene Name          | IL17RA   |
| Protein Name       | Interleukin-17 receptor A  |
| Immunogen          | The antiserum was produced against synthesized peptide derived from human IL17RA. AA range:141-190   |
| Specificity        | IL-17Rα Polyclonal Antibody detects endogenous levels of IL-17Rα protein.  |
| Formulation        | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  |
| Source             | Polyclonal, Rabbit,IgG   |
| Purification       | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  |
| Dilution           | Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not yet tested in other applications.  |
| Concentration      | 1 mg/ml  |
| Purity             | ≥90%   |
| Storage Stability  | -20°C/1 year   |
| Synonyms           | IL17RA; IL17R; Interleukin-17 receptor A; IL-17 receptor A; IL-17RA; CDw217; CD antigen CD217  |
| Observed Band      | 90kD   |
| Cell Pathway       | [Isoform 1]: Cell membrane ; Single-pass type I membrane protein .; [Isoform 2]:<br>Secreted .   |
| Tissue Specificity | Widely expressed.  |
| Function           | function:Receptor for IL17A. Binds its ligand with low affinity, suggesting that additional components are involved in IL17A-induced signaling.,PTM:Glycosylated.,tissue specificity:Widely expressed.,  |
| Background         | Interleukin 17A (IL17A) is a proinflammatory cytokine secreted by activated T-lymphocytes. It is a potent inducer of the maturation of CD34-positive hematopoietic precursors into neutrophils. The transmembrane protein encoded by this gene (interleukin 17A receptor; IL17RA) is a ubiquitous type I membrane glycoprotein that binds with low affinity to interleukin 17A. Interleukin 17A and its receptor play a pathogenic role in many inflammatory and autoimmune diseases |
|                    | Naniing Byahscience technology Co. Ltd   |

## Nanjing BYabscience technology Co.,Ltd



attention

国内优质抗体供应商

精准的 WB 检测服务 24H 在线服务, 欢迎咨询

such as rheumatoid arthritis. Like other cytokine receptors, this receptor likely has a multimeric structure. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by RefSeq, Feb 2014], Avoid repeated freezing and thawing! matters needing

Usage suggestions This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

| Products Images            |   |  |
|----------------------------|---|--|
| (kD)<br>117-<br>85- IL17RA | Western blot analysis of the lysates from HeLa cells using IL17RA antibody. |  |
| 48-                        |   |  |
| 34-                        |   |  |
| 26-                        |   |  |
| 19-                        |   |  |
| _                          |   |  |

Nanjing BYabscience technology Co., Ltd