




Olfactory receptor 10A7 Polyclonal Antibody

Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated		
Reactivity Human;Rat;Mouse; Applications WB;ELISA Gene Name OR10A7 Protein Name Olfactory receptor 10A7 Immunogen The antiserum was produced against synthesized peptide derived from human OR10A7. AA range:234-283 Specificity Olfactory receptor 10A7 Polyclonal Antibody detects endogenous levels of Olfactory receptor 10A7 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms OR10A7; Olfactory receptor 10A7; Olfactory receptor OR12-6 Observed Band 36kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,simillarity:Belongs to the G-protein coupled receptor family., Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-maily is the largest if transduction of odorant signals. The olfactory receptor family is the largest if transduction of odorant signals. The olfactory receptor family is the largest if transduction of odorant signals. The olfactory receptor genetic family is the largest if the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor genetic family is the largest if the protein family is the largest	Catalog No	BYab-13435
Applications WB;ELISA Gene Name OR10A7 Protein Name Olfactory receptor 10A7 Immunogen The antiserum was produced against synthesized peptide derived from human OR10A7. AA range:234-283 Specificity Olfactory receptor 10A7 Polyclonal Antibody detects endogenous levels of Olfactory receptor 10A7 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms OR10A7; Olfactory receptor 10A7; Olfactory receptor OR12-6 Observed Band 36kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor family., Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-maily is the largest if transduction of odorant signals. The olfactory receptor geneting is the largest in the protein family is the	Isotype	IgG
Gene Name OR10A7 Protein Name Olfactory receptor 10A7 Immunogen The antiserum was produced against synthesized peptide derived from human OR10A7. AA range:234-283 Specificity Olfactory receptor 10A7 Polyclonal Antibody detects endogenous levels of Olfactory receptor 10A7 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms OR10A7; Olfactory receptor 10A7; Olfactory receptor OR12-6 Observed Band 36kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor family., Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors share a -7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of dodrant signals. The olfactory receptor gene family is the largest it ransduction of dodrant signals. The olfactory receptor gene family is the largest it ransduction of dodrant signals. The olfactory receptor gene family is the largest it ransduction of dodrant signals. The olfactory receptor gene family is the largest it ransduction of dodrant signals. The olfactory receptor gene family is the largest it ransduction of dodrant signals. The olfactory receptor gene family is the largest it ransduction of dodrant signals. The olfactory receptor gene family is the largest it receptors.	Reactivity	Human;Rat;Mouse;
Protein Name Olfactory receptor 10A7 Immunogen The antiserum was produced against synthesized peptide derived from human OR10A7. AA range:234-283 Specificity Olfactory receptor 10A7 Polyclonal Antibody detects endogenous levels of Olfactory receptor 10A7 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms OR10A7; Olfactory receptor 10A7; Olfactory receptor OR12-6 Observed Band 36kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor family. Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of g-protein-coupled receptors (GPCR) arising from single codi	Applications	WB;ELISA
Immunogen The antiserum was produced against synthesized peptide derived from human OR10A7. AA range:234-283 Specificity Olfactory receptor 10A7 Polyclonal Antibody detects endogenous levels of Olfactory receptor 10A7 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications. Concentration 1 mg/ml Purity 290% Storage Stability -20°C/1 year Synonyms OR10A7; Olfactory receptor 10A7; Olfactory receptor OR12-6 Observed Band 36kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor family., Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors (GPCR) arising from single coding-exon genes. Olfactory receptors (GPCR) arising from single coding-exon genes. Olfactory receptors gene family is the largest it ransduction of odorant signals. The olfactory receptorine famility is the largest it ransduction of odorant signals. The olfactory receptorine famility is the largest it ransduction of odorant signals. The olfactory receptorine famility is the largest it ransduction of odorant signals. The olfactory receptorine family is the largest it ransduction of odorant signals. The olfactory receptorine family is the largest it ransduction of odorant signals. The olfactory receptorine family is the largest it ransduction of odorant signals. The olfactory receptor gene family is the largest it receptors.	Gene Name	OR10A7
OR10A7. AA range:234-283 Specificity Olfactory receptor 10A7 Polyclonal Antibody detects endogenous levels of Olfactory receptor 10A7 protein. Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms OR10A7; Olfactory receptor 10A7; Olfactory receptor OR12-6 Observed Band 36kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor family., Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest it ranget for odorant receptor gene family is the largest it ranget for odorant signals. The olfactory receptor gene family is the largest it ranget of odorant signals. The olfactory receptor gene family is the largest it range of odorant signals. The olfactory receptor gene family is the largest it ranges of odorant signals. The olfactory receptor gene family is the largest it ranges of odorant signals. The olfactory receptor gene family is the largest it ranges of odorant signals.	Protein Name	Olfactory receptor 10A7
Olfactory receptor 10A7 protein. Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms OR10A7; Olfactory receptor 10A7; Olfactory receptor OR12-6 Observed Band 36kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor family., Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory gene family is the largest it is lar	Immunogen	
Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms OR10A7; Olfactory receptor 10A7; Olfactory receptor OR12-6 Observed Band 36kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor family. Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest it has presented and protein family is the largest it is the largest it is protein.	Specificity	Olfactory receptor 10A7 Polyclonal Antibody detects endogenous levels of Olfactory receptor 10A7 protein.
Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms OR10A7; Olfactory receptor 10A7; Olfactory receptor OR12-6 Observed Band 36kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor family., Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of dodrant signals. The olfactory receptor gene family is the largest it has proved the following the following the following the following the following the following that the following t	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms OR10A7; Olfactory receptor 10A7; Olfactory receptor OR12-6 Observed Band 36kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor family., Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest it	Source	Polyclonal, Rabbit,IgG
applications. Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms OR10A7; Olfactory receptor 10A7; Olfactory receptor OR12-6 Observed Band 36kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor family., Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptor share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the nose of the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the nose of the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the nose of the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the nose of the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the nose of the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the nose of the receptor gene family is the largest in the nose of the receptor gene family is the largest in the nose of the receptor gene family is the largest in the nose of the receptor gene family is the largest in the nose of the receptor gene family is the largest in the nose of the receptor gene family is the largest in the nose of the receptor gene family is the largest in the nose of the receptor gene family is the largest in the nose of the receptor	Purification	
Purity ≥90% Storage Stability -20°C/1 year Synonyms OR10A7; Olfactory receptor 10A7; Olfactory receptor OR12-6 Observed Band 36kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor family., Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the signals.	Dilution	Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications.
Storage Stability -20°C/1 year Observed Band Gell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor family., Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptor (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the receptor gene family in the largest in the receptor gene family is the largest in the receptor gene family in the largest in the receptor gene family in the largest in t	Concentration	1 mg/ml
Synonyms OR10A7; Olfactory receptor 10A7; Olfactory receptor OR12-6 Observed Band 36kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor family., Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the largest in the receptor gene family is the largest in the l	Purity	≥90%
Observed Band Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor family., Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors have a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors have a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the nose, to initiate a neuronal response that triggers the perception of a smell.	Storage Stability	-20°C/1 year
Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor family., Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the nose, to initiate a neuronal response that triggers the perception of a smell.	Synonyms	OR10A7; Olfactory receptor 10A7; Olfactory receptor OR12-6
Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor family., Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a feet of the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a feet of the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the nose, to initiate a neuronal response that triggers the perception of a smell.	Observed Band	36kD
Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor family., Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the G-protein coupled receptor of family.	Cell Pathway	Cell membrane; Multi-pass membrane protein.
Background Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the nose, to initiate a neuronal receptors are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a receptors and are responsible for the recognition and G protein-mediated	Tissue Specificity	
neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest i	Function	function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family.,
	Background	neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658








	proteins for this organism is independent of other organisms. [provided by RefSeq, Jul 2008],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.





网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658