



IL-18R Polyclonal Antibody

Catalog No	BYab-13735
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB;ELISA
Gene Name	IL18R1
Protein Name	Interleukin-18 receptor 1
Immunogen	The antiserum was produced against synthesized peptide derived from the Internal region of human IL18R1. AA range:231-280
Specificity	IL-18R Polyclonal Antibody detects endogenous levels of IL-18R protein.
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	IL18R1; IL1RRP; Interleukin-18 receptor 1; IL-18R-1; IL-18R1; CD218 antigen-like family member A; CDw218a; IL1 receptor-related protein; IL-1Rrp; IL1R-rp; CD218a
Observed Band	63kD
Cell Pathway	Membrane ; Single-pass type I membrane protein .
Tissue Specificity	Highly expressed in leukocytes, spleen, lung. Also expressed, but at lower levels, in liver, small intestine, colon, prostate, thymus, placenta, and heart. Specifically coexpressed with IL18R1 in Th1 cells (PubMed:10925275, PubMed:11046021, PubMed:10653850).
Function	function:Receptor for interleukin 18 (IL-18). Binding to the agonist leads to the activation of NF-kappa-B.,miscellaneous:Does not bind interleukin 1 alpha (IL-1A) or interleukin 1 beta (IL-1B)..similarity:Belongs to the interleukin-1 receptor family.,similarity:Contains 1 TIR domain.,similarity:Contains 3 Ig-like C2-type (immunoglobulin-like) domains.,tissue specificity:Expressed in lung, leukocytes, spleen, liver, thymus, prostate, small intestine, colon, placenta, and heart, and is

Nanjing BYabscience technology Co.,Ltd



absent from brain, skeletal muscle, pancreas, and kidney. High level of expression in Hodgkin's disease cell lines.,

Background

The protein encoded by this gene is a cytokine receptor that belongs to the interleukin 1 receptor family. This receptor specifically binds interleukin 18 (IL18), and is essential for IL18 mediated signal transduction. IFN-alpha and IL12 are reported to induce the expression of this receptor in NK and T cells. This gene along with four other members of the interleukin 1 receptor family, including IL1R2, IL1R1, ILRL2 (IL-1Rrp2), and IL1RL1 (T1/ST2), form a gene cluster on chromosome 2q. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Sep 2013],


matters needing attention

Avoid repeated freezing and thawing!


Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western Blot analysis of 293 cells using IL-18R Polyclonal Antibody. Secondary antibody(catalog#:RS0002) was diluted at 1:20000



Western blot analysis of lysate from 293 cells, using IL18R1 Antibody.