



CD40 Monoclonal Antibody

Catalog No	BYab-13804
Isotype	IgG
Reactivity	Human
Applications	IF;ELISA
Gene Name	CD40
Protein Name	Tumor necrosis factor receptor superfamily member 5
Immunogen	Purified recombinant fragment of CD40 expressed in E. Coli.
Specificity	CD40 Monoclonal Antibody detects endogenous levels of CD40 protein.
Formulation	Purified recombinant fragment of CD40 expressed in E. Coli.
Source	Monoclonal, Mouse
Purification	Affinity purification
Dilution	Immunofluorescence: 1/200 - 1/1000. ELISA: 1/10000. Not yet tested in other applications.
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	CD40; TNFRSF5; Tumor necrosis factor receptor superfamily member 5; B-cell surface antigen CD40; Bp50; CD40L receptor; CDw40; CD antigen CD40
Observed Band	
Cell Pathway	[Isoform I]: Cell membrane; Single-pass type I membrane protein.; [Isoform II]: Secreted.
Tissue Specificity	B-cells and in primary carcinomas.
Function	alternative products:Additional isoforms seem to exist,disease:Defects in CD40 are the cause of hyper-IgM immunodeficiency type 3 (HIGM3) [MIM:606843]. HIGM3 is an autosomal recessive disorder which includes an inability of B cells to undergo isotype switching, one of the final differentiation steps in the humoral immune system, an inability to mount an antibody-specific immune response, and a lack of germinal center formation.,function:Receptor for TNFSF5/CD40LG.,online information:CD40 entry,online information:CD40 mutation db,similarity:Contains 4 TNFR-Cys repeats.,subunit:Monomer and homodimer. The variant form found in the bladder carcinoma cell line Hu549 does not form homodimers. Interacts with TRAF1, TRAF2, TRAF3, TRAF5 and TRAF6.,tissue specificity:B-cells and in primary carcinomas.,

Nanjing BYabscience technology Co.,Ltd



Background

This gene is a member of the TNF-receptor superfamily. The encoded protein is a receptor on antigen-presenting cells of the immune system and is essential for mediating a broad variety of immune and inflammatory responses including T cell-dependent immunoglobulin class switching, memory B cell development, and germinal center formation. AT-hook transcription factor AKNA is reported to coordinately regulate the expression of this receptor and its ligand, which may be important for homotypic cell interactions. Adaptor protein TNFR2 interacts with this receptor and serves as a mediator of the signal transduction. The interaction of this receptor and its ligand is found to be necessary for amyloid-beta-induced microglial activation, and thus is thought to be an early event in Alzheimer disease pathogenesis. Mutations affecting this gene are the cause of autosomal recessive hyper-IgM immunodeficiency type 3 (HIG


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Confocal immunofluorescence analysis of human peripheral blood lymphocytes (left) and mouse L1210 cells (right) using CD40 Monoclonal Antibody (green). Blue: DRAQ5 fluorescent DNA dye.